

Anti-IL-10 Antibody
Catalog # ABO12701**Specification**

Anti-IL-10 Antibody - Product Information

Application	WB, IHC-P, E
Primary Accession	P29456
Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Interleukin-10(IL10) detection. Tested with WB, IHC-P, ELISA in Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IL-10 Antibody - Additional Information

Gene ID 25325

Other Names

Interleukin-10, IL-10, Cytokine synthesis inhibitory factor, CSIF, IL10, IL-10

Calculated MW

20426 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Rat, By Heat

ELISA , 0.1-0.5 µg/ml, Rat, -
Western blot, 0.1-0.5 µg/ml, Rat

Subcellular Localization

Secreted.

Protein Name

Interleukin-10(IL-10)

Contents

Each vial contains 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃. Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify "Carrier Free" or "No BSA" in your order note. "

Immunogen

E. coli-derived rat IL-10 recombinant protein(Position: S19-N178).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-IL-10 Antibody - Protein Information

Name Il10

Synonyms Il-10

Function

Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3. In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators. Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro-inflammatory cytokines including granulocyte-macrophage colony-stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G-CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF-alpha. Also interferes with antigen presentation by reducing the expression of MHC- class II and co-stimulatory molecules, thereby inhibiting their ability to induce T cell activation (By similarity). In addition, controls the inflammatory response of macrophages by reprogramming essential metabolic pathways including mTOR signaling (By similarity).


Cellular Location

Secreted {ECO:0000250|UniProtKB:P22301}.

Anti-IL-10 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL-10 Antibody - Images



Anti-rat IL-10 antibody, ABO12701, Western blotting
Lane 1: Recombinant Rat IL-10 Protein 10ng
Lane 2: Recombinant Rat IL-10 Protein 5ng
Lane 3: Recombinant Rat IL-10 Protein 2

Anti-IL-10 Antibody - Background

Interleukin-10(IL-10 or IL10), also known as human cytokine synthesis inhibitory factor(CSIF), is an anti-inflammatory cytokine. In humans IL-10 is encoded by the IL10 gene. It is capable of inhibiting synthesis of pro-inflammatory cytokines like IFN-gamma, IL-2, IL-3, TNFalpha and GM-CSF made by cells such as macrophages and regulatory T-cells. IL-10 also displays potent abilities to suppress the antigen presentation capacity of antigen presenting cells. Kim et al.(1992) showed that the mouse IL 10 gene contains 5 exons and spans about 5.2 kb of genomic DNA. Eskdale et al.(1997) mapped the IL10 gene to the junction between 1q31 and 1q32.