

**Anti-FE65 Antibody**  
**Catalog # ABO12742****Specification**

---

**Anti-FE65 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O00213</a>
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Amyloid beta A4 precursor protein-binding family B member 1 (APBB1) detection. Tested with WB in Human;Mouse.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-FE65 Antibody - Additional Information**

**Gene ID** 322

**Other Names**

Amyloid beta A4 precursor protein-binding family B member 1, Protein Fe65, APBB1, FE65, RIR

**Calculated MW**

77244 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Mouse<br>

**Subcellular Localization**

Cell membrane. Cytoplasm. Nucleus. Cell projection, growth cone . Nucleus speckle. Colocalizes with TSHZ3 in axonal growth cone (By similarity). In normal conditions, it mainly localizes to the cytoplasm, while a small fraction is tethered to the cell membrane via its interaction with APP. Following exposure to DNA damaging agents, it is released from cell membrane and translocates to the nucleus. Nuclear translocation is under the regulation of APP. Colocalizes with TSHZ3 in the nucleus. Colocalizes with NEK6 at the nuclear speckles. Phosphorylation at Ser-610 by SGK1 promotes its localization to the nucleus (By similarity). .

**Tissue Specificity**

Highly expressed in brain; strongly reduced in post-mortem elderly subjects with Alzheimer disease. .

**Protein Name**

Amyloid beta A4 precursor protein-binding family B member 1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

**Immunogen**

E.coli-derived human FE65 recombinant protein (Position: Q295-A613). Human FE65 shares 95% and 96% amino acid (aa) sequence identity with mouse and rat FE65, respectively.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Contains 2 PID domains.

**Anti-FE65 Antibody - Protein Information**

**Name** APBB1 ([HGNC:581](#))

**Function**

Transcription coregulator that can have both coactivator and corepressor functions (PubMed: [15031292](http://www.uniprot.org/citations/15031292), PubMed: [18468999](http://www.uniprot.org/citations/18468999), PubMed: [18922798](http://www.uniprot.org/citations/18922798), PubMed: [25342469](http://www.uniprot.org/citations/25342469), PubMed: [33938178](http://www.uniprot.org/citations/33938178)). Adapter protein that forms a transcriptionally active complex with the gamma-secretase- derived amyloid precursor protein (APP) intracellular domain (PubMed: [15031292](http://www.uniprot.org/citations/15031292), PubMed: [18468999](http://www.uniprot.org/citations/18468999), PubMed: [18922798](http://www.uniprot.org/citations/18922798), PubMed: [25342469](http://www.uniprot.org/citations/25342469)). Plays a central role in the response to DNA damage by translocating to the nucleus and inducing apoptosis (PubMed: [15031292](http://www.uniprot.org/citations/15031292), PubMed: [18468999](http://www.uniprot.org/citations/18468999), PubMed: [18922798](http://www.uniprot.org/citations/18922798), PubMed: [25342469](http://www.uniprot.org/citations/25342469)). May act by specifically recognizing and binding histone H2AX phosphorylated on 'Tyr-142' (H2AXY142ph) at double-strand breaks (DSBs), recruiting other pro-apoptosis factors such as MAPK8/JNK1 (PubMed: [19234442](http://www.uniprot.org/citations/19234442), target=" \_blank">19234442</a>). Required for histone H4 acetylation at double-strand breaks (DSBs) (PubMed: [19234442](http://www.uniprot.org/citations/19234442), target=" \_blank">19234442</a>). Its ability to specifically bind modified histones and chromatin modifying enzymes such as KAT5/TIP60, probably explains its transcription activation activity (PubMed: [33938178](http://www.uniprot.org/citations/33938178), target=" \_blank">33938178</a>). Functions in association with TSHZ3, SET and HDAC factors as a transcriptional repressor, that inhibits the expression of CASP4 (PubMed: [19343227](http://www.uniprot.org/citations/19343227), target=" \_blank">19343227</a>). Associates with chromatin in a region surrounding the CASP4 transcriptional start site(s) (PubMed: [19343227](http://www.uniprot.org/citations/19343227), target=" \_blank">19343227</a>). Involved in hippocampal neurite branching and neuromuscular junction formation, as a result plays a role in spatial memory functioning (By similarity). Plays a role in the maintenance of lens transparency (By similarity). May play a role in muscle cell strength

(By similarity). Acts as a molecular adapter that functions in neurite outgrowth by activating the RAC1-ARF6 axis upon insulin treatment (PubMed:<a href="http://www.uniprot.org/citations/36250347" target="\_blank">36250347</a>).

#### Cellular Location

Cell membrane. Cytoplasm. Nucleus. Cell projection, growth cone {ECO:0000250|UniProtKB:P46933}. Nucleus speckle. Note=Colocalizes with TSHZ3 in axonal growth cone (By similarity). Colocalizes with TSHZ3 in the nucleus (PubMed:19343227). In normal conditions, it mainly localizes to the cytoplasm, while a small fraction is tethered to the cell membrane via its interaction with APP (PubMed:18468999). Following exposure to DNA damaging agents, it is released from cell membrane and translocates to the nucleus (PubMed:18468999). Nuclear translocation is under the regulation of APP (PubMed:18468999). Colocalizes with NEK6 at the nuclear speckles (PubMed:17512906). Phosphorylation at Ser-610 by SGK1 promotes its localization to the nucleus (By similarity) {ECO:0000250|UniProtKB:P46933, ECO:0000269|PubMed:17512906, ECO:0000269|PubMed:18468999, ECO:0000269|PubMed:19343227}

#### Tissue Location


Highly expressed in brain; strongly reduced in post-mortem elderly subjects with Alzheimer disease

#### Anti-FE65 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-FE65 Antibody - Images



Anti- FE65 antibody, ABO12742, Western blotting All lanes: Anti (ABO12742) at 0.5ug/mlWB:

Mouse Brain Tissue Lysate at 50ug Predicted bind size: 77KD Observed bind size: 77KD

### **Anti-FE65 Antibody - Background**

APBB1 is also known as RIR or FE65. The protein encoded by this gene is a member of the Fe65 protein family. It is an adaptor protein localized in the nucleus. It interacts with the Alzheimer's disease amyloid precursor protein (APP), transcription factor CP2/LSF/LBP1 and the low-density lipoprotein receptor-related protein. APP functions as a cytosolic anchoring site that can prevent the gene product's nuclear translocation. This encoded protein could play an important role in the pathogenesis of Alzheimer's disease. It is thought to regulate transcription. Also it is observed to block cell cycle progression by downregulating thymidylate synthase expression. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene.