

Anti-EAAT2 Antibody
Catalog # ABO12744**Specification**

Anti-EAAT2 Antibody - Product Information

Application	WB
Primary Accession	P43004
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Excitatory amino acid transporter 2(SLC1A2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-EAAT2 Antibody - Additional Information

Gene ID 6506

Other Names

Excitatory amino acid transporter 2, Glutamate/aspartate transporter II, Sodium-dependent glutamate/aspartate transporter 2, Solute carrier family 1 member 2, SLC1A2, EAAT2, GLT1

Calculated MW

62104 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Membrane; Multi-pass membrane protein.

Protein Name

Excitatory amino acid transporter 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived human EAAT2 recombinant protein (Position: T461-K574). Human EAAT2 shares 96% amino acid (aa) sequence identity with both mouse and rat EAAT2.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-EAAT2 Antibody - Protein Information

Name SLC1A2 ([HGNC:10940](#))

Function

Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed: [14506254](http://www.uniprot.org/citations/14506254), PubMed: [15265858](http://www.uniprot.org/citations/15265858), PubMed: [26690923](http://www.uniprot.org/citations/26690923), PubMed: [7521911](http://www.uniprot.org/citations/7521911)). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed: [14506254](http://www.uniprot.org/citations/14506254)). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed: [14506254](http://www.uniprot.org/citations/14506254)). Essential for the rapid removal of released glutamate from the synaptic cleft, and for terminating the postsynaptic action of glutamate (By similarity).

Cellular Location


Cell membrane; Multi-pass membrane protein


Anti-EAAT2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EAAT2 Antibody - Images





Anti-EAAT2 antibody, ABO12744, Western blotting All lanes: Anti EAAT2 (ABO12744) at 0.5ug/ml
Lane 1: Rat Brain Tissue Lysate at 50ug
Lane 2: Mouse Brain Tissue Lysate at 50ug
Lane 3: U87 Whole Cell Lysate at 40ug
Lane 4: SMMC Whole Cell Lysate at 40ug
Lane 5: PANC Whole Cell Lysate at 40ug
Lane 6: A549 Whole Cell Lysate at 40ug
Predicted bind size: 62KD
Observed bind size: 62KD

Anti-EAAT2 Antibody - Background

SLC1A2 is also known as EAAT2 or GLT-1. This gene encodes a member of a family of solute transporter proteins. The membrane-bound protein is the principal transporter that clears the excitatory neurotransmitter glutamate from the extracellular space at synapses in the central nervous system. Glutamate clearance is necessary for proper synaptic activation and to prevent neuronal damage from excessive activation of glutamate receptors. Mutations in and decreased expression of this protein are associated with amyotrophic lateral sclerosis. Alternatively spliced transcript variants of this gene have been identified.