

Anti-MAVS Picoband Antibody
Catalog # ABO12808**Specification****Anti-MAVS Picoband Antibody - Product Information**

Application	IHC
Primary Accession	Q7Z434
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Mitochondrial antiviral-signaling protein(MAVS) detection.
Tested with WB, IHC-P in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-MAVS Picoband Antibody - Additional Information**Gene ID 57506****Other Names**

Mitochondrial antiviral-signaling protein, MAVS, CARD adapter inducing interferon beta, Cardif, Interferon beta promoter stimulator protein 1, IPS-1, Putative NF-kappa-B-activating protein 031N, Virus-induced-signaling adapter, VISA, MAVS, IPS1, KIAA1271, VISA

Calculated MW

56528 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
 Western blot, 0.1-0.5 µg/ml, Human,

Subcellular Localization

Mitochondrion outer membrane. Mitochondrion. Peroxisome.

Tissue Specificity

Present in T-cells, monocytes, epithelial cells and hepatocytes (at protein level). Ubiquitously expressed, with highest levels in heart, skeletal muscle, liver, placenta and peripheral blood leukocytes. .

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E. coli-derived human MAVS recombinant protein (Position: L34-Q96). Human MAVS shares 72.6% and 69.4% amino acid (aa) sequence identity with mouse and rat MAVS, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-MAVS Picoband Antibody - Protein Information

Name MAVS {ECO:0000303|PubMed:16125763, ECO:0000312|HGNC:HGNC:29233}

Function

Adapter required for innate immune defense against viruses (PubMed:16125763, PubMed:16127453, PubMed:16153868, PubMed:16177806, PubMed:19631370, PubMed:20451243, PubMed:23087404, PubMed:20127681, PubMed:21170385, PubMed:27992402, PubMed:33139700, PubMed:37582970). Acts downstream of DHX33, RIGI and IFIH1/MDA5, which detect intracellular dsRNA produced during viral replication, to coordinate pathways leading to the activation of NF-kappa-B, IRF3 and IRF7, and to the subsequent induction of antiviral cytokines such as IFNB and RANTES (CCL5) (PubMed:16125763, PubMed:16127453, PubMed:16153868, PubMed:16177806, PubMed:19631370, PubMed:20451243, PubMed:23087404, PubMed:25636800, PubMed:20127681, PubMed:21170385, PubMed:20628368, PubMed:33110251, PubMed:27736772). Peroxisomal and mitochondrial MAVS act sequentially to create an antiviral cellular state (PubMed:20451243). Upon viral infection, peroxisomal MAVS induces the rapid interferon-independent expression of defense factors that provide short-term protection, whereas mitochondrial MAVS activates an interferon-dependent signaling pathway with delayed kinetics, which amplifies and stabilizes the antiviral response (PubMed:20451243). May activate the same pathways following detection of extracellular dsRNA by TLR3 (PubMed:16153868). May protect cells from apoptosis (PubMed:16125763). Involved in

NLRP3 inflammasome activation by mediating NLRP3 recruitment to mitochondria (PubMed:23582325).

Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Mitochondrion. Peroxisome

Tissue Location

Present in T-cells, monocytes, epithelial cells and hepatocytes (at protein level). Ubiquitously expressed, with highest levels in heart, skeletal muscle, liver, placenta and peripheral blood leukocytes.

Anti-MAVS Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Anti-MAVS Picoband Antibody - Images

Figure 5. IHC analysis of MAVS using anti-MAVS antibody (ABO12808).





Anti-MAVS Picoband Antibody - Background

Mitochondrial antiviral-signaling protein (MAVS) is a protein that in humans is encoded by the MAVS gene. The protein is also known by the names VISA (virus-induced signaling adapter), IPS-1 and Cardif. This gene encodes an intermediary protein necessary in the virus-triggered beta interferon signaling pathways. It is required for activation of transcription factors which regulate expression of beta interferon and contributes to antiviral immunity.