

# Anti-MMP13 Picoband Antibody

Catalog # ABO12834

Specification

### Anti-MMP13 Picoband Antibody - Product Information

ApplicationWB, IHC-P, EPrimary AccessionP23097HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for MMP13 detection. Tested with WB, IHC-P, Direct ELISA inHuman;Mouse;Rat.Human;Mouse;Rat.

**Reconstitution** Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

### Anti-MMP13 Picoband Antibody - Additional Information

Other Names Collagenase 3, 3.4.24.-, Matrix metalloproteinase-13, MMP-13, UMRCASE, Mmp13

**Application Details** Western blot, 0.1-0.5 μg/ml<br><br> Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/ml<br><br> Direct ELISA, 0.1-0.5 μg/ml<br>

**Subcellular Localization** Secreted, extracellular space, extracellular matrix. Secreted.

**Contents** Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

Immunogen E. coli-derived rat MMP13 recombinant protein (Position: Y99-H335).

**Cross Reactivity** No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and thawing.

# Anti-MMP13 Picoband Antibody - Protein Information



### Name Mmp13

#### Function

Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CCN2. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CCN2. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion (By similarity).

**Cellular Location** 

Secreted, extracellular space, extracellular matrix. Secreted

# Anti-MMP13 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### Anti-MMP13 Picoband Antibody - Images

<b>kDa</b> 180 - 130 -		
95 -		
72 -		
55 -	-	-
43 -		
34-	-	
26 -		
17-		

Figure 1. Western blot analysis of MMP13 using anti-MMP13 antibody (ABO12834). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat RH35 cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat



Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MMP13 antigen affinity purified polyclonal antibody (Catalog # ABO12834) at 0.5  $\hat{l}_{4}$ g/mL overnight at 4ŰC, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MMP13 at approximately 54KD. The expected band size for MMP13 is at 54KD.



Figure 2. IHC analysis of MMP13 using anti-MMP13 antibody (ABO12834).MMP13 was detected in paraffin-embedded section of mouse lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with  $1^{1}/4g/ml$  rabbit anti-MMP13 Antibody (ABO12834) overnight at 4ŰC. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37ŰC. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



Figure 3. IHC analysis of MMP13 using anti-MMP13 antibody (ABO12834).MMP13 was detected in paraffin-embedded section of mouse small intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with  $1\hat{l}_{4}$ g/ml rabbit anti-MMP13 Antibody (ABO12834) overnight at 4ŰC. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37ŰC. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.





Figure 4. IHC analysis of MMP13 using anti-MMP13 antibody (ABO12834).MMP13 was detected in paraffin-embedded section of rat lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with  $1^{1}/4$ g/ml rabbit anti-MMP13 Antibody (ABO12834) overnight at 4ŰC. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37ŰC. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



Figure 5. IHC analysis of MMP13 using anti-MMP13 antibody (ABO12834).MMP13 was detected in paraffin-embedded section of rat small intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with  $1\hat{l}_{4}^{1}$ g/ml rabbit anti-MMP13 Antibody (ABO12834) overnight at 4ŰC. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37ŰC. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

# Anti-MMP13 Picoband Antibody - Background

Collagenase 3 is an enzyme that in humans is encoded by the MMP13 gene. This gene encodes a member of the peptidase M10 family of matrix metalloproteinases (MMPs). Proteins in this family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. The encoded preproprotein is proteolytically processed to generate the mature protease. This protease cleaves type II collagen more efficiently than types I and III. It may be involved in articular cartilage turnover and cartilage pathophysiology associated with osteoarthritis. Mutations in this gene are associated with metaphyseal anadysplasia. This gene is part of a cluster of MMP genes on chromosome 11.