

Anti-IL17 Picoband Antibody
Catalog # ABO12835**Specification****Anti-IL17 Picoband Antibody - Product Information**

Application	WB, IHC-P, E
Primary Accession	Q62386
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for IL17 detection. Tested with WB, IHC-P, ELISA(Cap) in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IL17 Picoband Antibody - Additional Information

Gene ID 16171

Other Names

Interleukin-17A, IL-17, IL-17A, Cytotoxic T-lymphocyte-associated antigen 8, CTLA-8, Il17a, Ctla8, Il17

Application Details

Western blot, 0.1-0.5 µg/ml

 Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml

 ELISA(Cap), 0.1-0.5 µg/ml

Subcellular Localization

Secreted.

Tissue Specificity

Restricted to a subset of activated T-cells.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived mouse IL17 recombinant protein (Position: A26-A158).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r° Constitution, at 4°C; for one month. It° Can also be aliquotted and stored frozen at -20°C; for a

longer time. Avoid repeated freezing and thawing.

Anti-IL17 Picoband Antibody - Protein Information

Name Il17a

Synonyms Ctlα8, Il17

Function

Effector cytokine of innate and adaptive immune system involved in antimicrobial host defense and maintenance of tissue integrity (PubMed:18025225, PubMed:19144317, PubMed:26431948). Signals via IL17RA-IL17RC heterodimeric receptor complex, triggering homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter. This leads to downstream TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (PubMed:16200068, PubMed:17911633, PubMed:19144317, PubMed:26431948). Plays an important role in connecting T cell-mediated adaptive immunity and acute inflammatory response to destroy extracellular bacteria and fungi. As a signature effector cytokine of T-helper 17 cells (Th17), primarily induces neutrophil activation and recruitment at infection and inflammatory sites (PubMed:18025225). In airway epithelium, mediates neutrophil chemotaxis via induction of CXCL1 and CXCL5 chemokines (PubMed:18025225, PubMed:27923703). In secondary lymphoid organs, contributes to germinal center formation by regulating the chemotactic response of B cells to CXCL12 and CXCL13, enhancing retention of B cells within the germinal centers, B cell somatic hypermutation rate and selection toward plasma cells (PubMed:18157131). Effector cytokine of a subset of gamma-delta T cells that functions as part of an inflammatory circuit downstream IL1B, TLR2 and IL23A-IL12B to promote neutrophil recruitment for efficient bacterial clearance (PubMed:17372004, PubMed:20364087, PubMed:28709803). Effector cytokine of innate immune cells including invariant natural killer cell (iNKT) and group 3 innate lymphoid cells that mediate initial neutrophilic inflammation (PubMed:17470641, PubMed:23255360). Involved in the maintenance of the integrity of epithelial barriers during homeostasis and pathogen infection. Upon acute injury, has a direct role in epithelial barrier formation by regulating OCLN localization and tight junction biogenesis (PubMed:26431948). As part of the mucosal immune response induced by commensal bacteria, enhances host's ability to resist pathogenic bacterial and fungal infections by promoting neutrophil recruitment and antimicrobial peptides release (PubMed:28709803). In synergy with IL17F, mediates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (PubMed:19144317). Involved in antiviral host defense through various mechanisms (PubMed:21946434, PubMed:26735852, PubMed:27795421). Enhances immunity against West Nile virus by promoting T cell cytotoxicity (PubMed:27795421). May play a beneficial role in influenza A virus (H5N1) infection by enhancing B cell recruitment and immune response in the lung (PubMed:21946434). Contributes to influenza A virus (H1N1) clearance by driving the differentiation of B-1a B cells, providing for production of virus-specific IgM antibodies at first line of host defense (PubMed:26735852).

Cellular Location

Secreted.

Tissue Location

Expressed by Th17 cell lineage (at protein level). The expression pattern reflects the differentiation state, with IL17A- IL17F heterodimers produced at higher levels than IL17A-IL17A and IL17F-IL17F dimers in fully differentiated Th17 cells (PubMed:16990136, PubMed:18025225). Expressed in innate lymphoid cells (at protein level) (PubMed:23255360, PubMed:28709803). Expressed in gamma-delta T cell subsets (at protein level) (PubMed:17372004, PubMed:20364087, PubMed:26431948, PubMed:28709803). Expressed in iNKT cells (at protein level) (PubMed:17470641).

Anti-IL17 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL17 Picoband Antibody - Images




Figure 1. Western blot analysis of IL17 using anti-IL17 antibody (ABO12835). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human Jurkat cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL17 antigen affinity purified polyclonal antibody (Catalog # ABO12835) at 0.5 μ g/mL overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for IL17 at approximately 27KD. The expected band size for IL17 is at 17KD.






Figure 2. IHC analysis of IL17 using anti-IL17 antibody (ABO12835).



Anti-IL17 Picoband Antibody - Background

Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.