

Anti-EPO Receptor Picoband Antibody
Catalog # ABO12839**Specification**

Anti-EPO Receptor Picoband Antibody - Product Information

Application	WB, E
Primary Accession	P14753
Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for EPO Receptor detection. Tested with WB, Direct ELISA in Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-EPO Receptor Picoband Antibody - Additional Information

Gene ID 13857

Other Names

Erythropoietin receptor, EPO-R, Epor

Application Details

Western blot, 0.1-0.5 µg/ml

 Direct ELISA, 0.1-0.5 µg/ml

Subcellular Localization

Isoform EPOR-F: Cell membrane; Single-pass type I membrane protein.

Tissue Specificity

Expressed in relatively mature erythroid progenitor cells and in EPO-responsive erythroleukemia cells.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived mouse EPO Receptor recombinant protein (Position: D32-E225).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r° Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and

thawing.

Anti-EPO Receptor Picoband Antibody - Protein Information

Name Epor

Function

Receptor for erythropoietin. Mediates erythropoietin-induced erythroblast proliferation and differentiation. Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade. In some cell types, can also activate STAT1 and STAT3. May also activate the LYN tyrosine kinase.

Cellular Location

[Isoform EPOR-F]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in relatively mature erythroid progenitor cells and in EPO-responsive erythroleukemia cells

Anti-EPO Receptor Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EPO Receptor Picoband Antibody - Images

Anti-EPO Receptor Picoband Antibody - Background

The erythropoietin receptor (EpoR) is a protein that in humans is encoded by the EPOR gene. This gene encodes the erythropoietin receptor which is a member of the cytokine receptor family. Upon erythropoietin binding, this receptor activates Jak2 tyrosine kinase which activates different intracellular pathways including: Ras/MAP kinase, phosphatidylinositol 3-kinase and STAT transcription factors. The stimulated erythropoietin receptor appears to have a role in erythroid cell survival. Defects in the erythropoietin receptor may produce erythroleukemia and familial erythrocytosis. Dysregulation of this gene may affect the growth of certain tumors. Alternate splicing results in multiple transcript variants.