

Anti-TLS / FUS Picoband Antibody

Catalog # ABO12884

Specification

Anti-TLS / FUS Picoband Antibody - Product Information

ApplicationWBPrimary AccessionFUS: P35637HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for TLS / FUS detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-TLS / FUS Picoband Antibody - Additional Information

Application Details Western blot, 0.1-0.5 µg/ml

Subcellular Localization Nucleus

Tissue Specificity Ubiquitous.

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen A synthetic peptide corresponding to a sequence of human TLS / FUS (DNNTIFVQGLGENVTIESVADYFKQIGIIKTNKKT).

Purification Immunogen affinity purified.

Cross Reactivity No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-TLS / FUS Picoband Antibody - Protein Information



Anti-TLS / FUS Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-TLS / FUS Picoband Antibody - Images



Figure 1. Western blot analysis of TLS / FUS using anti-TLS / FUS antibody (ABO12884). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates,Lane 2: human K562 whole cell lysates,Lane 3: mouse NIH3T3 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TLS / FUS antigen affinity purified polyclonal antibody (Catalog # ABO12884) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for TLS / FUS at approximately 75KD. The expected band size for TLS / FUS is at 53KD.

Anti-TLS / FUS Picoband Antibody - Background

RNA-binding protein FUS/TLS (Fused in Sarcoma/Translocated in Sarcoma) is a protein that in humans is encoded by the FUS gene. This gene encodes a multifunctional protein component of the heterogeneous nuclear ribonucleoprotein (hnRNP) complex. The hnRNP complex is involved in pre-mRNA splicing and the export of fully processed mRNA to the cytoplasm. This protein belongs to the FET family of RNA-binding proteins which have been implicated in cellular processes that include regulation of gene expression, maintenance of genomic integrity and mRNA/microRNA



processing. Alternative splicing results in multiple transcript variants. Defects in this gene result in amyotrophic lateral sclerosis type 6.