

## Anti-Psoriasin Picoband Antibody

Catalog # ABO12982

#### Specification

## Anti-Psoriasin Picoband Antibody - Product Information

ApplicationWB, IHC-P, EPrimary AccessionP31151HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Psoriasin detection. Tested with WB, IHC-P, Direct ELISA inHuman; Mouse; Rat.

**Reconstitution** Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

### Anti-Psoriasin Picoband Antibody - Additional Information

Gene ID 6278

**Other Names** Protein S100-A7, Psoriasin, S100 calcium-binding protein A7, S100A7, PSOR1, S100A7C

Calculated MW 11471 MW KDa

**Application Details** Western blot, 0.1-0.5 μg/ml<br><br> Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/ml<br><br> Direct ELISA, 0.1-0.5 μg/ml<br>

**Subcellular Localization** Cytoplasm . Secreted . Secreted by a non- classical secretory pathway.

Tissue Specificity

Fetal ear, skin, and tongue and human cell lines. Highly up-regulated in psoriatic epidermis. Also highly expressed in the urine of bladder squamous cell carcinoma (SCC) bearing patients.

**Contents** Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

Immunogen E. coli-derived human Psoriasin recombinant protein (Position: S2-Q101).

**Cross Reactivity** No cross reactivity with other proteins.



Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and thawing.

# Anti-Psoriasin Picoband Antibody - Protein Information

Name S100A7

Synonyms PSOR1, S100A7C

**Cellular Location** Cytoplasm. Secreted. Note=Secreted by a non-classical secretory pathway

**Tissue Location** 

Fetal ear, skin, and tongue and human cell lines. Highly up-regulated in psoriatic epidermis. Also highly expressed in the urine of bladder squamous cell carcinoma (SCC) bearing patients

### **Anti-Psoriasin Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### **Anti-Psoriasin Picoband Antibody - Images**



Figure 1. Western blot analysis of Psoriasin using anti-Psoriasin antibody (ABO12982). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing



conditions. Lane 1: human placenta tissue lysates, Lane 2: human Hela cell lysates, Lane 3: rat spleen tissue lysates, Lane 4: rat RH35 cell lysates, Lane 5: mouse spleen tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Psoriasin antigen affinity purified polyclonal antibody (Catalog # ABO12982) at 0.5  $\hat{l}$ /4g/mL overnight at 4ŰC, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Psoriasin at approximately 17KD. The expected band size for Psoriasin is at 11KD.



Figure 2. IHC analysis of Psoriasin using anti-Psoriasin antibody (ABO12982).Psoriasin was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with  $1\hat{l}_{4}$ g/ml rabbit anti-Psoriasin Antibody (ABO12982) overnight at 4ŰC. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37ŰC. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



Figure 3. IHC analysis of Psoriasin using anti-Psoriasin antibody (ABO12982).Psoriasin was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with  $1\hat{l}_{4}$ g/ml rabbit anti-Psoriasin Antibody (ABO12982) overnight at 4ŰC. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37ŰC. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

# Anti-Psoriasin Picoband Antibody - Background

S100 calcium-binding protein A7 (S100A7), also known as psoriasin, is a protein that in humans is encoded by the S100A7 gene. The protein encoded by this gene is a member of the S100 family of



proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein differs from the other S100 proteins of known structure in its lack of calcium binding ability in one EF-hand at the N-terminus. The protein is overexpressed in hyperproliferative skin diseases, exhibits antimicrobial activities against bacteria and induces immunomodulatory activities.