

### **Anti-Myf5 Picoband Antibody**

Catalog # ABO13024

## **Specification**

# **Anti-Myf5 Picoband Antibody - Product Information**

Application WB, E
Primary Accession P13349
Host Rabbit

Reactivity Human, Mouse, Rat

Clonality Polyclonal Lyophilized

**Description** 

Rabbit IgG polyclonal antibody for Myf5 detection. Tested with WB, Direct ELISA in

Human; Mouse; Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

### **Anti-Myf5 Picoband Antibody - Additional Information**

**Gene ID 4617** 

**Other Names** 

Myogenic factor 5, Myf-5, Class C basic helix-loop-helix protein 2, bHLHc2, MYF5, BHLHC2

**Application Details** 

Western blot, 0.1-0.5 μg/ml<br> onless ELISA, 0.1-0.5 μg/ml<br/> br>

**Contents** 

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immuno**aen

E. coli-derived human Myf5 recombinant protein (Position: E25-Y146).

**Cross Reactivity** 

No cross reactivity with other proteins.

Storage At -20°C; for one year. After r°Constitution,

at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and

thawing.

## **Anti-Myf5 Picoband Antibody - Protein Information**

Name MYF5



#### **Synonyms** BHLHC2

#### **Function**

Transcriptional activator that promotes transcription of muscle-specific target genes and plays a role in muscle differentiation (PubMed:<a href="http://www.uniprot.org/citations/29887215" target="\_blank">29887215</a>). Together with MYOG and MYOD1, co-occupies muscle- specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into myoblasts. Probable sequence specific DNA-binding protein.

**Cellular Location** Nucleus.

### **Anti-Myf5 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### **Anti-Myf5 Picoband Antibody - Images**





Figure 1. Western blot analysis of Myf5 using anti-Myf5 antibody (ABO13024).

#### **Anti-Myf5 Picoband Antibody - Background**

Myogenic factor 5 is a protein that in humans is encoded by the MYF5 gene. It is a protein with a key role in regulating muscle differentiation or myogenesis, specifically the development of skeletal muscle. Myf5 belongs to a family of proteins known as myogenic regulatory factors (MRFs). This transcription factor is the earliest of all MRFs to be expressed in the embryo, where it is only markedly expressed for a few days (specifically around 8 days post-somite formation and lasting until day 14 post-somite in mice). It functions during that time to commit myogenic precursor cells





to become skeletal muscle. In fact, its expression in proliferating myoblasts has led to its classification as a determination factor. Furthermore, Myf5 is a master regulator of muscle development, possessing the ability to induce a muscle phenotype upon its forced expression in fibroblastic cells.