

## **Anti-LAMP3 Picoband Antibody**

Catalog # ABO13068

## **Specification**

# **Anti-LAMP3 Picoband Antibody - Product Information**

Application IHC
Primary Accession Q9UQV4
Host Rabbit

Reactivity Human, Mouse, Rat

Clonality Polyclonal Lyophilized

**Description** 

Rabbit IgG polyclonal antibody for LAMP3 detection. Tested with IHC-P, Direct ELISA in Human; Mouse; Rat.

#### Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

## **Anti-LAMP3 Picoband Antibody - Additional Information**

#### **Gene ID 27074**

#### **Other Names**

Lysosome-associated membrane glycoprotein 3, LAMP-3, Lysosomal-associated membrane protein 3, DC-lysosome-associated membrane glycoprotein, DC LAMP, Protein TSC403, CD208, LAMP3, DCLAMP, TSC403

### **Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1  $\mu$ g/ml<br/>br> Direct ELISA, 0.1-0.5  $\mu$ g/ml<br/>br>

# **Subcellular Localization**

Lysosome membrane.

#### **Tissue Specificity**

Detected in tonsil interdigitating dendritic cells, in spleen, lymph node, Peyer's patches in the small instestine, in thymus medulla and in B-cells (at protein level). Expressed in lymphoid organs and dendritic cells. Expressed in lung. Up-regulated in carcinomas of the esophagus, colon, rectum, ureter, stomach, breast, fallopian tube, thyroid and parotid tissues.

#### **Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

### **Immunogen**

E. coli-derived human LAMP3 recombinant protein (Position: K28-E375).

## **Cross Reactivity**

No cross reactivity with other proteins.



Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and thawing.

## **Anti-LAMP3 Picoband Antibody - Protein Information**

Name LAMP3

Synonyms DCLAMP, TSC403

#### **Function**

Lysosomal membrane glycoprotein which plays a role in the unfolded protein response (UPR) that contributes to protein degradation and cell survival during proteasomal dysfunction (PubMed:<a href="http://www.uniprot.org/citations/25681212" target="\_blank">25681212</a>). Plays a role in the process of fusion of the lysosome with the autophagosome, thereby modulating the autophagic process (PubMed:<a href="http://www.uniprot.org/citations/24434718" target="\_blank">24434718</a>). Promotes hepatocellular lipogenesis through activation of the PI3K/Akt pathway (PubMed:<a href="http://www.uniprot.org/citations/29056532" target="\_blank">29056532</a>). May also play a role in dendritic cell function and in adaptive immunity (PubMed:<a href="http://www.uniprot.org/citations/9768752" target=" blank">9768752</a>).

#### **Cellular Location**

Cell surface. Lysosome membrane; Single-pass type I membrane protein. Cytoplasmic vesicle membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Note=During dendritic cell maturation, detected on cytoplasmic vesicles (the MHC II compartment) that contain MHC II proteins, LAMP1, LAMP2 and LAMP3 (PubMed:9768752). Detected on lysosomes in mature dendritic cells (PubMed:9768752).

#### **Tissue Location**

Detected in tonsil interdigitating dendritic cells, in spleen, lymph node, Peyer's patches in the small instestine, in thymus medulla and in B-cells (at protein level). Expressed in lymphoid organs and dendritic cells. Expressed in lung. Up-regulated in carcinomas of the esophagus, colon, rectum, ureter, stomach, breast, fallopian tube, thyroid and parotid tissues

## **Anti-LAMP3 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Anti-LAMP3 Picoband Antibody - Images






Figure 4. IHC analysis of LAMP3 using anti-LAMP3 antibody (ABO13068).

## **Anti-LAMP3 Picoband Antibody - Background**

Lysosome-associated membrane glycoprotein 3 (LAMP3, Lamp3) is a protein that in humans is encoded by theLAMP3 gene. This gene is mapped to chromosome 3q26.3-q27. It is one of the lysosome-associated membrane glycoproteins. Dendritic cells (DCs) are the most potent antigen-presenting cells. Immature DCs efficiently capture antigens and differentiate into interdigitating dendritic cells (IDCs) in lymphoid tissues that induce primary T-cell responses