

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody
Catalog # ABO13193

Specification

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	Q99717
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody - Additional Information

Gene ID 4090

Other Names

Mothers against decapentaplegic homolog 5, MAD homolog 5, Mothers against DPP homolog 5, JV5-1, SMAD family member 5, SMAD 5, Smad5, hSmad5, SMAD5, MADH5

Calculated MW

52258 MW KDa

Application Details

WB 1:500-1:1000

Subcellular Localization

Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.

Tissue Specificity

Ubiquitous.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Phospho-Smad1/5/9

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody - Protein Information

Name SMAD5 ([HGNC:6771](#))

Synonyms MADH5

Function

Transcriptional regulator that plays a role in various cellular processes including embryonic development, cell differentiation, angiogenesis and tissue homeostasis (PubMed:[12064918](http://www.uniprot.org/citations/12064918), PubMed:[16516194](http://www.uniprot.org/citations/16516194)). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRI) and associates with SMAD4 to form a heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed:[9442019](http://www.uniprot.org/citations/9442019)). In turn, the hetero-trimeric complex recognizes cis- regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:[33510867](http://www.uniprot.org/citations/33510867)). Non-phosphorylated SMAD5 has a cytoplasmic role in energy metabolism regulation by promoting mitochondrial respiration and glycolysis in response to cytoplasmic pH changes (PubMed:[28675158](http://www.uniprot.org/citations/28675158)). Mechanistically, interacts with hexokinase 1/HK1 and thereby accelerates glycolysis (PubMed:[28675158](http://www.uniprot.org/citations/28675158)).

Cellular Location

Cytoplasm. Nucleus Mitochondrion. Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4


Tissue Location


Ubiquitous.

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Phospho-Smad1/5/9 (S463/S465/S467) Rabbit Monoclonal Antibody - Images



Western blot analysis of Phospho-Smad5 (S463/S465) expression in (1) HeLa cell lysate; (2) HeLa cell treated with BMP-4 lysate.