

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody
Catalog # ABO13485**Specification**

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC
Primary Accession	P04179
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody - Additional Information

Gene ID 6648

Other Names

Superoxide dismutase [Mn], mitochondrial, 1.15.1.1, SOD2

Calculated MW

24722 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200

Subcellular Localization

Mitochondrion matrix.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human SOD2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody - Protein Information

Name SOD2

Function

Destroys superoxide anion radicals which are normally produced within the cells and which are toxic to biological systems.

Cellular Location

Mitochondrion matrix.

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SOD2/Mnsod Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of TOMM20/Tom20 using anti-TOMM20/Tom20 antibody (M04039). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,
Lane 2: human 293T whole cell lysates,
Lane 3: human K562 whole cell lysates,
Lane 4: human HeLa whole cell lysates,
Lane 5: rat kidney tissue lysates,
Lane 6: rat stomach tissue lysates,
Lane 7: mouse kidney tissue lysates,
Lane 8: mouse stomach tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was

incubated with rabbit anti-TOMM20/Tom20 antigen affinity purified monoclonal antibody (Catalog # M04039) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TOMM20/Tom20 at approximately 16 kDa. The expected band size for TOMM20/Tom20 is at 16 kDa.