

**Anti-CD80/B7 1 Rabbit Monoclonal Antibody**  
Catalog # ABO13507

**Specification**

---

**Anti-CD80/B7 1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P33681</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-CD80/B7 1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

**Anti-CD80/B7 1 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 941

**Other Names**

T-lymphocyte activation antigen CD80, Activation B7-1 antigen, BB1, CTLA-4 counter-receptor B7.1, B7, CD80, CD80, CD28LG, CD28LG1, LAB7

**Calculated MW**

33048 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200

**Subcellular Localization**

Membrane; Single-pass type I membrane protein.

**Tissue Specificity**

Expressed on activated B-cells, macrophages and dendritic cells.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human CD80

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated**

## freeze-thaw cycles.

### Anti-CD80/B7 1 Rabbit Monoclonal Antibody - Protein Information

**Name** CD80

**Synonyms** CD28LG, CD28LG1, LAB7

#### Function

Costimulatory molecule that belongs to the immunoglobulin superfamily that plays an important role in T-lymphocyte activation (PubMed:<a href="http://www.uniprot.org/citations/38467718" target="\_blank">38467718</a>). Acts as the primary auxiliary signal augmenting the MHC/TCR signal in naive T-cells together with the CD28 receptor which is constitutively expressed on the cell surface of T-cells (PubMed:<a href="http://www.uniprot.org/citations/12196291" target="\_blank">12196291</a>). In turn, activates different signaling pathways such as NF-kappa-B or MAPK leading to the production of different cytokines (PubMed:<a href="http://www.uniprot.org/citations/10438913" target="\_blank">10438913</a>). In addition, CD28/CD80 costimulatory signal stimulates glucose metabolism and ATP synthesis of T-cells by activating the PI3K/Akt signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/12121659" target="\_blank">12121659</a>). Also acts as a regulator of PDL1/PDCD1 interactions to limit excess engagement of PDL1 and its inhibitory role in immune responses (PubMed:<a href="http://www.uniprot.org/citations/36727298" target="\_blank">36727298</a>). Expressed on B-cells, plays a critical role in regulating interactions between B-cells and T-cells in both early and late germinal center responses, which are crucial for the generation of effective humoral immune responses (By similarity).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Expressed on activated B-cells, macrophages and dendritic cells


### Anti-CD80/B7 1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-CD80/B7 1 Rabbit Monoclonal Antibody - Images





All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of CD80 expression in Raji cell lysate.