

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody

Catalog # ABO13608

Specification

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody - Product Information

Application WB, IHC, IF, ICC, IP, FC

Primary Accession
Host
Rabbit
Isotype
Rabbit IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

Description

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody - Additional Information

Gene ID 4323

Other Names

Matrix metalloproteinase-14, MMP-14, 3.4.24.80, MMP-X1, Membrane-type matrix metalloproteinase 1, MT-MMP 1, MTMMP1, Membrane-type-1 matrix metalloproteinase, MT1-MMP, MT1MMP, MMP14

Calculated MW

65894 MW KDa

Application Details

WB 1:1000-1:2000
br>IHC 1:50-1:200
br>ICC/IF 1:50-1:200
br>IP 1:50
br>FC 1:50

Subcellular Localization

Membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Forms a complex with BST2 and localizes to the cytoplasm.

Tissue Specificity

Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors...

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human MMP14

Purification

Affinity-chromatography



Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody - Protein Information

Name MMP14

Function

Endopeptidase that degrades various components of the extracellular matrix such as collagen (PubMed:8015608). Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development (By similarity). Activates progelatinase A/MMP2, thereby acting as a positive regulator of cell growth and migration (PubMed: 22065321, PubMed:8015608). Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed: 12714657, PubMed:22065321). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:20837484). Acts as a regulator of Notch signaling by mediating cleavage and inhibition of DLL1 (PubMed:21572390). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:22330140). Acts as a negative regulator of the GDF15-GFRAL aversive response by mediating cleavage and inactivation of GFRAL (PubMed:35177851).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Forms a complex with BST2 and localizes to the cytoplasm (PubMed:17081065)

Tissue Location

Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.


Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody - Images





Western blot analysis of MMP14 expression in human spleen lysate.



Immunohistochemical analysis of paraffin-embedded human kdieny, using MMP14 Antibody.