

**Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody**  
**Catalog # ABO13728****Specification****Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">Q06830</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.

**Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody - Additional Information****Gene ID** 5052**Other Names**

Peroxiredoxin-1, 1.11.1.24, Natural killer cell-enhancing factor A, NKEF-A, Proliferation-associated gene protein, PAG, Thioredoxin peroxidase 2, Thioredoxin-dependent peroxide reductase 2, Thioredoxin-dependent peroxiredoxin 1, PRDX1, PAGA, PAGB, TDpx2

**Calculated MW**

22110 MW KDa

**Application Details**

WB 1:1000-1:2000&lt;br&gt;IHC 1:50-1:200&lt;br&gt;ICC/IF 1:50-1:200&lt;br&gt;FC 1:50

**Subcellular Localization**

Cytoplasm. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human PRDX1

**Purification**

Affinity-chromatography

**Storage**

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

## Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody - Protein Information

**Name** PRDX1

**Synonyms** PAGA, PAGB, TDPX2

### Function

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H<sub>2</sub>O<sub>2</sub> (PubMed:<a href="http://www.uniprot.org/citations/9497357" target="\_blank">9497357</a>). Reduces an intramolecular disulfide bond in GPD5 that gates the ability to GPD5 to drive postmitotic motor neuron differentiation (By similarity).

### Cellular Location


Cytoplasm. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

## Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-PRDX1/Peroxiredoxin 1 Rabbit Monoclonal Antibody - Images



Western blot analysis of PRDX1 expression in K562 cell lysates.



All lanes use the Antibody at 1:3W dilution for 1 hour at room temperature.