

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody

Catalog # ABO13784

Specification

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody - Product Information

Application Primary Accession Host Isotype Reactivity Clonality Format Description WB, IHC, IF, ICC, IP, FC <u>013148</u> Rabbit Rabbit IgG Rat, Human, Mouse Monoclonal Liquid

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody - Additional Information

Gene ID 23435

Other Names TAR DNA-binding protein 43, TDP-43, TARDBP {ECO:0000303|PubMed:18396105, ECO:0000312|HGNC:HGNC:11571}

Calculated MW 44740 MW KDa

Application Details WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:200

Subcellular Localization Nucleus. In patients with frontotemporal lobar degeneration and amyotrophic lateral sclerosis, it is absent from the nucleus of affected neurons but it is the primary component of cytoplasmic ubiquitin-positive inclusion bodies.

Tissue Specificity Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract and spleen.

Contents Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen A synthesized peptide derived from human TDP43

Purification Affinity-chromatography



Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody - Protein Information

Name TARDBP {ECO:0000303|PubMed:18396105, ECO:0000312|HGNC:HGNC:11571}

Function

RNA-binding protein that is involved in various steps of RNA biogenesis and processing (PubMed:23519609). Preferentially binds, via its two RNA recognition motifs RRM1 and RRM2, to GU-repeats on RNA molecules predominantly localized within long introns and in the 3'UTR of mRNAs (PubMed:23519609, PubMed:24240615, PubMed:24464995). In turn, regulates the splicing of many non-coding and protein-coding RNAs including proteins involved in neuronal survival, as well as mRNAs that encode proteins relevant for neurodegenerative diseases (PubMed:21358640, PubMed:29438978). Plays a role in maintaining mitochondrial homeostasis by regulating the processing of mitochondrial transcripts (PubMed:28794432). Also regulates mRNA stability by recruiting CNOT7/CAF1 deadenylase on mRNA 3'UTR leading to poly(A) tail deadenylation and thus shortening (PubMed:30520513). In response to oxidative insult, associates with stalled ribosomes localized to stress granules (SGs) and contributes to cell survival (PubMed: 19765185, PubMed:23398327). Also participates in the normal skeletal muscle formation and regeneration, forming cytoplasmic myo-granules and binding mRNAs that encode sarcomeric proteins (PubMed:30464263). Plays a role in the maintenance of the circadian clock periodicity via stabilization of the CRY1 and CRY2 proteins in a FBXL3-dependent manner (PubMed:27123980). Negatively regulates the expression of CDK6 (PubMed:19760257). Regulates the expression of HDAC6, ATG7 and VCP in a PPIA/CYPA-dependent manner (PubMed:25678563).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, Stress granule Mitochondrion. Note=Continuously travels in and out of the nucleus (PubMed:18957508). Localizes to stress granules in response to oxidative stress (PubMed:19765185). A small subset localizes in mitochondria (PubMed:28794432).

Tissue Location

Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract and spleen

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Western Blot</u>



- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-TDP43 TARDBP Rabbit Monoclonal Antibody - Images



All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.





Western blot analysis of TDP43 expression in K562 cell lysate.