

Anti-RALBP1 Rabbit Monoclonal Antibody
Catalog # ABO13838**Specification****Anti-RALBP1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, FC
Primary Accession	Q15311
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-RALBP1 Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-RALBP1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 10928

Other Names

RalA-binding protein 1, 76 kDa Ral-interacting protein, Dinitrophenyl S-glutathione ATPase, DNP-SG ATPase, 7.6.2.2, 7.6.2.3, Ral-interacting protein 1, RALBP1 ([HGNC:9841](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=9841))

Calculated MW

76063 MW KDa

Application Details

WB 1:5000-1:10000
IHC 1:50-1:200
FC 1:50

Subcellular Localization

Membrane ; Peripheral membrane protein.

Tissue Specificity

Expressed ubiquitously but at low levels. Shows a strong expression in the erythrocytes..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human RALBP1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-RALBP1 Rabbit Monoclonal Antibody - Protein Information

Name RALBP1 ([HGNC:9841](#))

Function

Multifunctional protein that functions as a downstream effector of RALA and RALB (PubMed:7673236). As a GTPase-activating protein/GAP can inactivate CDC42 and RAC1 by stimulating their GTPase activity (PubMed:7673236). As part of the Ral signaling pathway, may also regulate ligand-dependent EGF and insulin receptors-mediated endocytosis (PubMed:10910768, PubMed:12775724). During mitosis, may act as a scaffold protein in the phosphorylation of EPSIN/EPN1 by the mitotic kinase cyclin B-CDK1, preventing endocytosis during that phase of the cell cycle (PubMed:12775724). During mitosis, also controls mitochondrial fission as an effector of RALA (PubMed:21822277). Recruited to mitochondrion by RALA, acts as a scaffold to foster the mitotic kinase cyclin B-CDK1-mediated phosphorylation and activation of DNMI1 (PubMed:21822277).

Cellular Location

Cell membrane; Peripheral membrane protein. Cytoplasm, cytosol Cytoplasm, cytoskeleton, spindle pole {ECO:0000250|UniProtKB:Q62796} Nucleus. Mitochondrion. Note=Cytosolic protein that transiently associates with the mitotic spindle poles in early prophase, and dissociates from them after completion of mitosis (By similarity) Targeted to the plasma membrane through its interaction with RALB, directed by FGF signaling. Docking on the membrane is required to transduce the Ral signal (By similarity). Recruited by RALA to the mitochondrion during mitosis where it regulates mitochondrial fission (PubMed:21822277). Nuclear localization is cell cycle dependent while membrane localization is seen in adherent cells (PubMed:22319010). The region involved in membrane association could form transmembrane domains and expose a part of the protein extracellularly (Probable) {ECO:0000250|UniProtKB:Q62796, ECO:0000250|UniProtKB:Q9PT60, ECO:0000269|PubMed:21822277, ECO:0000269|PubMed:22319010, ECO:0000305|PubMed:15610018}

Tissue Location

Expressed ubiquitously but at low levels. Shows a strong expression in the erythrocytes.


Anti-RALBP1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Anti-RALBP1 Rabbit Monoclonal Antibody - Images



Western blot analysis of RALBP1 expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded human breast, using RALBP1 Antibody.