

Anti-Moesin Rabbit Monoclonal Antibody
Catalog # ABO13841**Specification****Anti-Moesin Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P26038
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Moesin Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-Moesin Rabbit Monoclonal Antibody - Additional Information

Gene ID 4478

Other Names

Moesin, Membrane-organizing extension spike protein, MSN (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=7373)
HGNC:7373

Calculated MW

67820 MW KDa

Application Details

WB 1:1000-1:2000
IHC 1:50-1:100
ICC/IF 1:50-1:100
IP 1:20
FC 1:20

Subcellular Localization

Cell membrane ; Peripheral membrane protein ; Cytoplasmic side. Cytoplasm, cytoskeleton. Apical cell membrane ; Peripheral membrane protein ; Cytoplasmic side. Cell projection, microvillus membrane ; Peripheral membrane protein ; Cytoplasmic side. Phosphorylated form is enriched in microvilli-like structures at apical membrane (By similarity). Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment..

Tissue Specificity

In all tissues and cultured cells studied.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Moesin

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Moesin Rabbit Monoclonal Antibody - Protein Information

Name MSN ([HGNC:7373](#))

Function

Ezrin-radixin-moesin (ERM) family protein that connects the actin cytoskeleton to the plasma membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (PubMed:10212266). Once phosphorylated on its C-terminal threonine, moesin is activated leading to interaction with F-actin and cytoskeletal rearrangement (PubMed:10212266). These rearrangements regulate many cellular processes, including cell shape determination, membrane transport, and signal transduction (PubMed:12387735, PubMed:15039356). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:9298994, PubMed:9616160). Modulates phagolysosomal biogenesis in macrophages (By similarity). Also participates in immunologic synapse formation (PubMed:27405666).

Cellular Location

Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P26041}. Apical cell membrane {ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus membrane {ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus {ECO:0000250|UniProtKB:P26041}. Note=Phosphorylated form is enriched in microvilli-like structures at apical membrane. Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment (By similarity). Localizes at the uropods of T lymphoblasts. {ECO:0000250|UniProtKB:P26041, ECO:0000269|PubMed:18586956, ECO:0000269|PubMed:9298994}

Tissue Location

In all tissues and cultured cells studied.


Anti-Moesin Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Moesin Rabbit Monoclonal Antibody - Images



Western blot analysis of Moesin expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded human thyroid cancer, using Moesin Antibody.