

## Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody

**Catalog # ABO13986** 

#### **Specification**

# **Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody - Product Information**

Application WB, IHC, IF, ICC, IP

Primary Accession

Host
Isotype
Reactivity
Clonality
Format

P55211
Rabbit
Rabbit IgG
Human, Mouse
Monoclonal
Liquid

**Description** 

Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse.

# Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody - Additional Information

#### Gene ID 842

#### **Other Names**

Caspase-9, CASP-9, 3.4.22.62, Apoptotic protease Mch-6, Apoptotic protease-activating factor 3, APAF-3, ICE-like apoptotic protease 6, ICE-LAP6, Caspase-9 subunit p35, Caspase-9 subunit p10, CASP9, MCH6

# Calculated MW 46281 MW KDa

# **Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50</br>

#### **Tissue Specificity**

Ubiquitous, with highest expression in the heart, moderate expression in liver, skeletal muscle, and pancreas. Low levels in all other tissues. Within the heart, specifically expressed in myocytes...

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### **Immunogen**

A synthesized peptide derived from human Caspase-9

# **Purification**

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.



# Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody - Protein Information

Name CASP9

**Synonyms MCH6** 

#### **Function**

Involved in the activation cascade of caspases responsible for apoptosis execution. Binding of caspase-9 to Apaf-1 leads to activation of the protease which then cleaves and activates effector caspases caspase-3 (CASP3) or caspase-7 (CASP7). Promotes DNA damage- induced apoptosis in a ABL1/c-Abl-dependent manner. Proteolytically cleaves poly(ADP-ribose) polymerase (PARP). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed:<a href="http://www.uniprot.org/citations/36758105" target="\_blank">36758105</a>, PubMed:<a href="http://www.uniprot.org/citations/36758106" target="\_blank">36758106</a>).

#### **Tissue Location**


Ubiquitous, with highest expression in the heart, moderate expression in liver, skeletal muscle, and pancreas. Low levels in all other tissues. Within the heart, specifically expressed in myocytes.

# **Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Anti-Caspase-9 CASP9 Rabbit Monoclonal Antibody - Images



Immunohistochemical analysis of paraffin-embedded human liver, using Caspase-9 Antibody.





Immunofluorescent analysis of HeLa cells treated with staurosporine, using Caspase-9 Antibody.




Figure 1. Western blot analysis of Caspase-9 using anti-Caspase-9 antibody (M00080). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human K562 whole cell lysates,

Lane 3: human SiHa whole cell lysates,

Lane 4: human HEL whole cell lysates,

Lane 5: human Jurkat whole cell lysates,

Lane 6: human lurkat whole cell lysates,

Lane 7: mouse NIH/3T3 whole cell lysates,

Lane 8: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Caspase-9 antigen affinity purified monoclonal antibody (Catalog # M00080) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Caspase-9 at approximately 46 kDa. The expected band size for Caspase-9 is at 46 kDa.