

Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody
Catalog # ABO13994**Specification****Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P46734/P52564
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody - Additional Information**Calculated MW**

39318 MW KDa

Application Details

WB 1:500-1:1000
IHC 1:50-1:200
ICC/IF 1:100-1:500
IP 1:50
FC 1:50

Tissue Specificity

Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human MEK3/MEK6

Purification

Affinity-chromatography

Storage


Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody - Protein Information**Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-MEK3/MEK6 MAP2K3 Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.



Western blot analysis of MEK3/MEK6 expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded mouse spleen, using MEK3/MEK6 Antibody.



Immunofluorescent analysis of HeLa cells, using MEK3/MEK6 Antibody.