

Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody

Catalog # ABO14212

Specification

Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody - Product Information

Application WB, IHC, IF, ICC, FC

Primary Accession
Host
Rabbit
Isotype
Rabbit IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

Description

Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 7001

Other Names

Peroxiredoxin-2, 1.11.1.24, Natural killer cell-enhancing factor B, NKEF-B, PRP, Thiol-specific antioxidant protein, TSA, Thioredoxin peroxidase 1, Thioredoxin-dependent peroxide reductase 1, Thioredoxin-dependent peroxiredoxin 2, PRDX2, NKEFB, TDPX1

Calculated MW

21892 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50</br>

Subcellular Localization

Cytoplasm.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Peroxiredoxin 2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.



Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody - Protein Information

Name PRDX2

Synonyms NKEFB, TDPX1

Function

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).


Cellular Location Cytoplasm.

Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Peroxiredoxin 2 Rabbit Monoclonal Antibody - Images



Western blot analysis of Peroxiredoxin 2 expression in HEK293 cell lysate.