

**Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody**  
**Catalog # ABO14213****Specification****Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">P05164</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications.  
This antibody reacts with Human.

**Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 4353

**Other Names**

Myeloperoxidase, MPO, 1.11.2.2, Myeloperoxidase, 89 kDa myeloperoxidase, 84 kDa myeloperoxidase, Myeloperoxidase light chain, Myeloperoxidase heavy chain, MPO ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=7218](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=7218))  
HGNC:7218

**Calculated MW**

83869 MW KDa

**Application Details**

WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200

**Subcellular Localization**

Lysosome.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Myeloperoxidase

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody - Protein Information

**Name** MPO ([HGNC:7218](#))

### Function

Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity (PubMed:<a href="http://www.uniprot.org/citations/9922160" target="\_blank">9922160</a>). Mediates the proteolytic cleavage of alpha-1-microglobulin to form t-alpha-1-microglobulin, which potently inhibits oxidation of low-density lipoprotein particles and limits vascular damage (PubMed:<a href="http://www.uniprot.org/citations/25698971" target="\_blank">25698971</a>).

### Cellular Location

Lysosome.

## Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of MPO using anti-MPO antibody (M00372).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HL-60 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MPO antigen affinity purified monoclonal antibody (Catalog # M00372) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MPO at approximately 84 kDa. The expected band size for MPO is at 84 kDa.