

**Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody**  
**Catalog # ABO14219****Specification**

---

**Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP
Primary Accession	<a href="#">Q13509</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 10381

**Other Names**

Tubulin beta-3 chain, Tubulin beta-4 chain, Tubulin beta-III, TUBB3, TUBB4

**Calculated MW**

50433 MW KDa

**Application Details**

WB 1:2000-1:20000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50

**Subcellular Localization**

Cytoplasm, cytoskeleton.

**Tissue Specificity**

Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues..

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human beta III Tubulin

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated**

## freeze-thaw cycles.

### Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody - Protein Information

**Name** TUBB3

**Synonyms** TUBB4

#### Function

Tubulin is the major constituent of microtubules, protein filaments consisting of alpha- and beta-tubulin heterodimers (PubMed: [34996871](http://www.uniprot.org/citations/34996871), PubMed: [38305685](http://www.uniprot.org/citations/38305685), PubMed: [38609661](http://www.uniprot.org/citations/38609661)). Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms (PubMed: [34996871](http://www.uniprot.org/citations/34996871), PubMed: [38305685](http://www.uniprot.org/citations/38305685), PubMed: [38609661](http://www.uniprot.org/citations/38609661)). Below the cap, alpha-beta tubulin heterodimers are in GDP-bound state, owing to GTPase activity of alpha-tubulin (PubMed: [34996871](http://www.uniprot.org/citations/34996871), PubMed: [38609661](http://www.uniprot.org/citations/38609661)). TUBB3 plays a critical role in proper axon guidance and maintenance (PubMed: [20074521](http://www.uniprot.org/citations/20074521)). Binding of NTN1/Netrin-1 to its receptor UNC5C might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed: [28483977](http://www.uniprot.org/citations/28483977)). Plays a role in dorsal root ganglion axon projection towards the spinal cord (PubMed: [28483977](http://www.uniprot.org/citations/28483977)).

#### Cellular Location

Cytoplasm, cytoskeleton. Cell projection, growth cone {ECO:0000250|UniProtKB:Q9ERD7}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q9ERD7}. Cell projection, filopodium {ECO:0000250|UniProtKB:Q9ERD7}

#### Tissue Location


Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues.

### Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-beta III Tubulin TUBB3 Rabbit Monoclonal Antibody - Images



Immunohistochemical analysis of paraffin-embedded human ovarian cancer, using beta III Tubulin Antibody(M01857)




TUBB3 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-TUBB3 Antibody (M01857)overnight at 4 Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.





Western blot analysis of beta III Tubulin expression in HeLa cell lysate (M01857).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.


After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TUBB3 monoclonal antibody (Catalog # M01857) overnight at 4 then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TUBB3







AB0042Immunofluorescent analysis of PC12 cells, using beta Tubulin Antibody.



Immunofluorescent analysis using the Antibody at 1:50 dilution.