

**Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody**  
Catalog # ABO14263**Specification****Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q8NEB9</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

**Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 5289

**Other Names**

Phosphatidylinositol 3-kinase catalytic subunit type 3, PI3-kinase type 3, PI3K type 3, PtdIns-3-kinase type 3, 2.7.1.137, Phosphatidylinositol 3-kinase p100 subunit, Phosphoinositide-3-kinase class 3, hVps34, PIK3C3 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=8974](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=8974) target="\_blank">HGNC:8974</a>), VPS34 {ECO:0000305}

**Calculated MW**

101549 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200

**Subcellular Localization**

Midbody. Late endosome. Cytoplasmic vesicle, autophagosome. As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes (Probable). Localizes also to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme (By similarity)..

**Tissue Specificity**

Ubiquitously expressed, with a highest expression in skeletal muscle..

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human PI 3 Kinase Class 3

**Purification**

Affinity-chromatography

**Storage****Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.****Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody - Protein Information****Name** PIK3C3 ([HGNC:8974](#))**Synonyms** VPS34 {ECO:0000305}**Function**

Catalytic subunit of the PI3K complex that mediates formation of phosphatidylinositol 3-phosphate; different complex forms are believed to play a role in multiple membrane trafficking pathways: PI3KC3-C1 is involved in initiation of autophagosomes and PI3KC3-C2 in maturation of autophagosomes and endocytosis (PubMed: [14617358](http://www.uniprot.org/citations/14617358), PubMed: [33637724](http://www.uniprot.org/citations/33637724), PubMed: [7628435](http://www.uniprot.org/citations/7628435)). As part of PI3KC3-C1, promotes endoplasmic reticulum membrane curvature formation prior to vesicle budding (PubMed: [32690950](http://www.uniprot.org/citations/32690950)). Involved in regulation of degradative endocytic trafficking and required for the abscission step in cytokinesis, probably in the context of PI3KC3-C2 (PubMed: [20208530](http://www.uniprot.org/citations/20208530), PubMed: [20643123](http://www.uniprot.org/citations/20643123)). Involved in the transport of lysosomal enzyme precursors to lysosomes (By similarity). Required for transport from early to late endosomes (By similarity).

**Cellular Location**

Midbody. Late endosome. Cytoplasmic vesicle, autophagosome. Note=As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes (PubMed:14617358). Also localizes to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme (By similarity) {ECO:0000250|UniProtKB:Q6PF93, ECO:0000305|PubMed:14617358}

**Tissue Location**


Ubiquitously expressed, with a highest expression in skeletal muscle.

**Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-PI 3 Kinase Class 3 PIK3C3 Rabbit Monoclonal Antibody - Images**



Western blot analysis of PI 3 Kinase Class 3 expression in 293T cell lysate.



Immunohistochemical analysis of paraffin-embedded human kidney, using PI 3 Kinase Class 3 Antibody.