

**Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10)**  
**Catalog # ABO14347****Specification**

---

**Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) - Product Information**

Application	WB, IHC, ICC, FC
Primary Accession	<a href="#">P50402</a>
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) . Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) - Additional Information**

**Gene ID** 2010

**Other Names**

Emerin, EMD, EDMD, STA

**Calculated MW**

34 kDa KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml<br> Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml<br> Immunohistochemistry (Frozen Section), 0.5-1 µg/ml<br> Immunocytochemistry, 0.5-1 µg/ml<br> Flow Cytometry, 1-3 µg/1x10<sup>6</sup> cells<br>

**Subcellular Localization**

Nucleus inner membrane

**Tissue Specificity**

Skeletal muscle, heart, colon, testis, ovary and pancreas.

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>N.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human Emerin, different from the related mouse sequence by eight amino acids, and from the related rat sequence by nine amino acids.

**Cross Reactivity**

No cross-reactivity with other proteins.

Storage

**Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.**

## **Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) - Protein Information**

**Name** EMD

**Synonyms** EDMD, STA

### **Function**

Stabilizes and promotes the formation of a nuclear actin cortical network. Stimulates actin polymerization in vitro by binding and stabilizing the pointed end of growing filaments. Inhibits beta- catenin activity by preventing its accumulation in the nucleus. Acts by influencing the nuclear accumulation of beta-catenin through a CRM1- dependent export pathway. Links centrosomes to the nuclear envelope via a microtubule association. Required for proper localization of non- farnesylated prelamin-A/C. Together with NEMP1, contributes to nuclear envelope stiffness in germ cells (PubMed:<a href="http://www.uniprot.org/citations/32923640" target="\_blank">32923640</a>). EMD and BAF are cooperative cofactors of HIV-1 infection. Association of EMD with the viral DNA requires the presence of BAF and viral integrase. The association of viral DNA with chromatin requires the presence of BAF and EMD.

### **Cellular Location**

Nucleus inner membrane; Single-pass membrane protein; Nucleoplasmic side. Nucleus outer membrane. Note=Colocalized with BANF1 at the central region of the assembling nuclear rim, near spindle-attachment sites. The accumulation of different intermediates of prelamin-A/C (non-farnesylated or carboxymethylated farnesylated prelamin-A/C) in fibroblasts modify its localization in the nucleus

### **Tissue Location**


Skeletal muscle, heart, colon, testis, ovary and pancreas

## **Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) - Images**






Figure 1. Western blot analysis of Emerin using anti-Emerin antibody (M00714).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human placenta tissue lysates,

Lane 3: human Caco-2 whole cell lysates,

Lane 4: human HepG2 whole cell lysates,

Lane 5: Rabbit IgG,

Lane 6: Marker 1113,

Lane 7: human Jurkat whole cell lysates.

Lane 8: human MDA-MB-453 whole cell lysates,

Lane 9: human SK-OV-3 whole cell lysates,

Lane 10: human SW620 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Emerin antigen affinity purified monoclonal antibody (Catalog # M00714) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.




Figure 3. IHC analysis of Emerin using anti-Emerin antibody (M00714).

Emerin was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Emerin Antibody (M00714) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1021) with DAB as the

chromogen.




Figure 2. IHC analysis of Emerin using anti-Emerin antibody (M00714).

Emerin was detected in paraffin-embedded section of human gastric cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu\text{g}/\text{ml}$  mouse anti-Emerin Antibody (M00714) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1021) with DAB as the chromogen.




Figure 4. Flow Cytometry analysis of A431 cells using anti-Emerin antibody (M00714).

Overlay histogram showing A431 cells stained with M00714 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Emerin Antibody (M00714, 1  $\mu\text{g}/1 \times 10^6$  cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10  $\mu\text{g}/1 \times 10^6$  cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1  $\mu\text{g}/1 \times 10^6$ ) used under the same conditions. Unlabelled sample (Red line) was also used as a control.