

Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody
Catalog # ABO14368**Specification****Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P45983/P45984/P53779
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody - Additional Information**Application Details**

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Phospho-JNK1/2/3 (T183+T183+T221)
Phospho-JNK1/2/3 (T183+T183+T221)

Purification

Affinity-chromatography

Storage


Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody - Protein Information**Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-Phospho-JNK1/2/3 (T183+T183+T221) MAPK8 Monoclonal Antibody - Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




Immunofluorescent analysis using the Antibody at 1:50 dilution.




Immunofluorescent analysis using the Antibody at 1:50 dilution.



Western blot analysis of JNK1/2/3 phosphorylation expression in NIH/3T3 cell lysate treated with Anisomycin.



Immunohistochemical analysis of paraffin-embedded human breast, using Phospho-JNK1/2/3 (T183+T183+T221) Antibody.



Immunofluorescent analysis of NIH/3T3 cells treated with Anisomycin, using Phospho-JNK1/2/3 (T183+T183+T221) Antibody.