

**Anti-CD82 Monoclonal Antibody**  
**Catalog # ABO14489****Specification**

---

**Anti-CD82 Monoclonal Antibody - Product Information**

Application	WB, IHC, FC
Primary Accession	<a href="#">P27701</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-CD82 Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human, Rat.

**Anti-CD82 Monoclonal Antibody - Additional Information**

**Gene ID** 3732

**Other Names**

CD82 antigen, C33 antigen, IA4, Inducible membrane protein R2, Metastasis suppressor Kangai-1, Suppressor of tumorigenicity 6 protein, Tetraspanin-27, Tspan-27, CD82, CD82, KAI1, SAR2, ST6, TSPAN27

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>FC 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human CD82 Associates with CD4 or CD8 and delivers costimulatory signals for the TCR/CD3 pathway.

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-CD82 Monoclonal Antibody - Protein Information**

**Name** CD82

**Synonyms** KAI1, SAR2, ST6, TSPAN27**Function**

Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling (PubMed:<a href="http://www.uniprot.org/citations/19497983" target="\_blank">19497983</a>). Participates thereby in diverse biological functions such as cell signal transduction, adhesion, migration and protein trafficking. Acts as a attenuator of EGF signaling, facilitating ligand-induced endocytosis of the receptor and its subsequent desensitization (PubMed:<a href="http://www.uniprot.org/citations/10985391" target="\_blank">10985391</a>, PubMed:<a href="http://www.uniprot.org/citations/35538033" target="\_blank">35538033</a>). Mechanistically, modulates ligand- induced ubiquitination and trafficking of EGFR via E3 ligase CBL phosphorylation by PKC (PubMed:<a href="http://www.uniprot.org/citations/23897813" target="\_blank">23897813</a>). Increases cell-matrix adhesion by regulating the membrane organization of integrin alpha4/ITA4 (PubMed:<a href="http://www.uniprot.org/citations/24623721" target="\_blank">24623721</a>, PubMed:<a href="http://www.uniprot.org/citations/8757325" target="\_blank">8757325</a>). Modulates adhesion and suppresses cell migration through other integrins such as the alpha6/ITGA6 and beta1/ITGB1 (PubMed:<a href="http://www.uniprot.org/citations/15557282" target="\_blank">15557282</a>, PubMed:<a href="http://www.uniprot.org/citations/17560548" target="\_blank">17560548</a>). Decreases cell-associated plasminogen activation by interfering with the interaction between urokinase-type plasminogen activator/PLAU and its receptor PLAU (PubMed:<a href="http://www.uniprot.org/citations/15677461" target="\_blank">15677461</a>). Associates with CD4 or CD8 and delivers costimulatory signals for the TCR/CD3 pathway. Plays a role in TLR9 trafficking to acidified CpG-containing compartments by controlling interaction between TLR9 and VAMP3 and subsequent myddosome assembly (By similarity). Inhibits LPS-induced inflammatory response by preventing binding of LPS to TLR4 on the cell surface (PubMed:<a href="http://www.uniprot.org/citations/36945827" target="\_blank">36945827</a>). Plays a role in the activation of macrophages into anti-inflammatory phenotypes (By similarity). Independently of Toll- like receptor (TLR) signaling, is recruited to pathogen-containing phagosomes prior to fusion with lysosomes and thereby participates in antigen presentation (By similarity). Also acts to control angiogenesis and switch angiogenic milieu to quiescent state by binding and sequestering VEGFA and PDGFB to inhibit the signaling they trigger via their respective cell surface receptor (PubMed:<a href="http://www.uniprot.org/citations/34530889" target="\_blank">34530889</a>).

**Cellular Location**

Cell membrane {ECO:0000269|PubMed:19497983, ECO:0000269|PubMed:23897813, ECO:0000269|PubMed:30463011, ECO:0000269|PubMed:34530889, ECO:0000269|PubMed:8757325, ECO:0000269|Ref.4}; Multi-pass membrane protein Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:P40237}

**Tissue Location**

Lymphoid specific.


**Anti-CD82 Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-CD82 Monoclonal Antibody - Images



Western blot analysis of CD82 expression in Jurkat cell lysate.