

**Anti-TRPM8 Monoclonal Antibody**  
**Catalog # ABO14516****Specification**

---

**Anti-TRPM8 Monoclonal Antibody - Product Information**

Application	WB, IF, ICC
Primary Accession	<a href="#">Q7Z2W7</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-TRPM8 Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse.

**Anti-TRPM8 Monoclonal Antibody - Additional Information**

**Gene ID** 79054

**Other Names**

Transient receptor potential cation channel subfamily M member 8, Long transient receptor potential channel 6, LTrpC-6, LTrpC6, Transient receptor potential p8, Trp-p8, TRPM8, LTRPC6, TRPP8

**Application Details**

WB 1:500-1:2000<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human TRPM8

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-TRPM8 Monoclonal Antibody - Protein Information**

**Name** TRPM8

**Synonyms** LTRPC6, TRPP8

### Function

Non-selective ion channel permeable to monovalent and divalent cations, including Na(+), K(+), and Ca(2+), with higher permeability for Ca(2+). Activated by multiple factors, such as temperature, voltage, pressure, and changes in osmolality. Activated by cool temperatures (<23-28 degrees Celsius) and by chemical ligands evoking a sensation of coolness, such as menthol and icilin therefore plays a central role in the detection of environmental cold temperatures (PubMed:<a href="http://www.uniprot.org/citations/15306801" target="\_blank">15306801</a>, PubMed:<a href="http://www.uniprot.org/citations/15852009" target="\_blank">15852009</a>, PubMed:<a href="http://www.uniprot.org/citations/16174775" target="\_blank">16174775</a>, PubMed:<a href="http://www.uniprot.org/citations/25559186" target="\_blank">25559186</a>, PubMed:<a href="http://www.uniprot.org/citations/37857704" target="\_blank">37857704</a>). TRPM8 is a voltage-dependent channel; its activation by cold or chemical ligands shifts its voltage thresholds towards physiological membrane potentials, leading to the opening of the channel (PubMed:<a href="http://www.uniprot.org/citations/15306801" target="\_blank">15306801</a>). In addition to its critical role in temperature sensing, regulates basal tear secretion by sensing evaporation-induced cooling and changes in osmolality (By similarity). May plays a role in prostate cancer cell migration (PubMed:<a href="http://www.uniprot.org/citations/16174775" target="\_blank">16174775</a>, PubMed:<a href="http://www.uniprot.org/citations/25559186" target="\_blank">25559186</a>).

### Cellular Location

Cell membrane; Multi-pass membrane protein. Membrane raft {ECO:0000250|UniProtKB:Q8R4D5}. Endoplasmic reticulum membrane. Note=Lipid raft association modulates TRPM8 channel activity (By similarity) Located in the endoplasmic reticulum in prostate cancer cells (PubMed:11325849, PubMed:16174775). {ECO:0000250|UniProtKB:Q8R4D5, ECO:0000269|PubMed:11325849, ECO:0000269|PubMed:16174775}

### Tissue Location


Expressed in prostate. Also expressed in prostate tumors and in non-prostatic primary tumors such as colon, lung, breast and skin tumors.


## Anti-TRPM8 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-TRPM8 Monoclonal Antibody - Images





All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Western blot analysis of TRPM8 expression in A549 cell lysate.

### **Anti-TRPM8 Monoclonal Antibody - Background**

Activated by icilin, eucalyptol, menthol, cold and modulation of intracellular pH. Involved in menthol sensation. Permeable for monovalent cations sodium, potassium, and cesium and divalent cation calcium.