

Anti-CD97 Monoclonal Antibody

Catalog # ABO14543

Specification

# Anti-CD97 Monoclonal Antibody - Product Information

Application Primary Accession Host Isotype Reactivity Clonality Format Description WB, IHC, IF, ICC, IP, FC <u>P48960</u> Rabbit Rabbit IgG Human, Mouse Monoclonal Liquid

Anti-CD97 Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse.

# Anti-CD97 Monoclonal Antibody - Additional Information

Gene ID 976

Other Names

Adhesion G protein-coupled receptor E5 {ECO:0000312|HGNC:HGNC:1711}, Leukocyte antigen CD97, CD97, Adhesion G protein-coupled receptor E5 subunit alpha, Adhesion G protein-coupled receptor E5 subunit beta, ADGRE5 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=1711" target="\_blank">HGNC:1711</a>)

Application Details WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50<br>FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CD97 Receptor potentially involved in both adhesion and signaling processes early after leukocyte activation. Plays an essential role in leukocyte migration.

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

### Anti-CD97 Monoclonal Antibody - Protein Information



# Name ADGRE5 (HGNC:1711)

### **Function**

Receptor potentially involved in both adhesion and signaling processes early after leukocyte activation. Plays an essential role in leukocyte migration.

Cellular Location Cell membrane {ECO:0000250|UniProtKB:Q9Z0M6}; Multi-pass membrane protein

#### Tissue Location

Broadly expressed, found on most hematopoietic cells, including activated lymphocytes, monocytes, macrophages, dendritic cells, and granulocytes. Expressed also abundantly by smooth muscle cells. Expressed in thyroid, colorectal, gastric, esophageal and pancreatic carcinomas too. Expression are increased under inflammatory conditions in the CNS of multiple sclerosis and in synovial tissue of patients with rheumatoid arthritis. Increased expression of CD97 in the synovium is accompanied by detectable levels of soluble CD97 in the synovial fluid

### Anti-CD97 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### Anti-CD97 Monoclonal Antibody - Images



Western blot analysis of CD97 expression in U937 cell lysate.