

**Anti-Ctip1 Monoclonal Antibody**  
**Catalog # ABO14550****Specification****Anti-Ctip1 Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">Q9H165</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Ctip1 Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Ctip1 Monoclonal Antibody - Additional Information**

**Gene ID** 53335

**Other Names**

B-cell lymphoma/leukemia 11A, BCL-11A, B-cell CLL/lymphoma 11A, COUP-TF-interacting protein 1, Ecotropic viral integration site 9 protein homolog, EVI-9, Zinc finger protein 856, BCL11A, CTIP1, EVI9, KIAA1809, ZNF856

**Application Details**

WB 1:1000-1:5000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:100

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Ctip1 Functions as a myeloid and B-cell proto-oncogene. May play important roles in leukemogenesis and hematopoiesis. An essential factor in lymphopoiesis, is required for B-cell formation in fetal liver. May function as a modulator of the transcriptional repression activity of ARP1.

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Ctip1 Monoclonal Antibody - Protein Information**

**Name** BCL11A

**Synonyms** CTIP1, EVI9, KIAA1809, ZNF856

### Function

Transcription factor (PubMed:<a href="http://www.uniprot.org/citations/16704730" target="\_blank">16704730</a>, PubMed:<a href="http://www.uniprot.org/citations/29606353" target="\_blank">29606353</a>). Associated with the BAF SWI/SNF chromatin remodeling complex (PubMed:<a href="http://www.uniprot.org/citations/23644491" target="\_blank">23644491</a>, PubMed:<a href="http://www.uniprot.org/citations/39607926" target="\_blank">39607926</a>). Binds to the 5'-TGACCA-3' sequence motif in regulatory regions of target genes, including a distal promoter of the HBG1 hemoglobin subunit gamma-1 gene (PubMed:<a href="http://www.uniprot.org/citations/29606353" target="\_blank">29606353</a>, PubMed:<a href="http://www.uniprot.org/citations/39423807" target="\_blank">39423807</a>). Involved in regulation of the developmental switch from gamma- to beta-globin, probably via direct repression of HBG1; hence indirectly repressing fetal hemoglobin (HbF) level (PubMed:<a href="http://www.uniprot.org/citations/26375765" target="\_blank">26375765</a>, PubMed:<a href="http://www.uniprot.org/citations/29606353" target="\_blank">29606353</a>, PubMed:<a href="http://www.uniprot.org/citations/39423807" target="\_blank">39423807</a>, PubMed:<a href="http://www.uniprot.org/citations/39607926" target="\_blank">39607926</a>). Involved in brain development (PubMed:<a href="http://www.uniprot.org/citations/27453576" target="\_blank">27453576</a>). May play a role in hematopoiesis (By similarity). Essential factor in lymphopoiesis required for B-cell formation in fetal liver (By similarity). May function as a modulator of the transcriptional repression activity of NR2F2 (By similarity).

### Cellular Location

Cytoplasm. Nucleus. Chromosome. Note=Associates with the nuclear body Colocalizes with SUMO1 and SENP2 in nuclear speckles (By similarity) [Isoform 2]: Cytoplasm. Nucleus Note=Predominantly localized in the nucleus in nuclear paraspeckles

### Tissue Location


Expressed at high levels in brain, spleen thymus, bone marrow and testis. Expressed in CD34-positive myeloid precursor cells, B-cells, monocytes and megakaryocytes. Expression is tightly regulated during B-cell development.


### Anti-Ctip1 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Ctip1 Monoclonal Antibody - Images





Western blot analysis of CtIp1 expression in Jurkat cell lysate.




Figure 1. Western blot analysis of CtIp1 using anti-CtIp1 antibody (M00741).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CtIp1 antigen affinity purified monoclonal antibody (Catalog # M00741) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CtIp1 at approximately 120 kDa. The expected band size for CtIp1 is at 84 kDa.