

**Anti-KPNA2 Monoclonal Antibody**  
**Catalog # ABO14603****Specification****Anti-KPNA2 Monoclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, FC                 |
| Primary Accession | <a href="#">P52292</a> |
| Host              | Rabbit                 |
| Isotype           | Rabbit IgG             |
| Reactivity        | Rat, Human, Mouse      |
| Clonality         | Monoclonal             |
| Format            | Liquid                 |

**Description**

Anti-KPNA2 Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-KPNA2 Monoclonal Antibody - Additional Information**

**Gene ID** 3838

**Other Names**

Importin subunit alpha-1, Karyopherin subunit alpha-2, RAG cohort protein 1, SRP1-alpha, KPNA2 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=6395](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6395)), RCH1, SRP1

**Application Details**

WB 1:1000-1:5000<br>FC 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human KPNA2 Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. Binds specifically and directly to substrates containing either a simple or bipartite NLS motif.

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-KPNA2 Monoclonal Antibody - Protein Information**

**Name** KPNA2 ([HGNC:6395](#))

**Synonyms** RCH1, SRP1**Function**

Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1 (PubMed:<a href="http://www.uniprot.org/citations/28991411" target="\_blank">28991411</a>, PubMed:<a href="http://www.uniprot.org/citations/32130408" target="\_blank">32130408</a>, PubMed:<a href="http://www.uniprot.org/citations/7604027" target="\_blank">7604027</a>, PubMed:<a href="http://www.uniprot.org/citations/7754385" target="\_blank">7754385</a>). Binds specifically and directly to substrates containing either a simple or bipartite NLS motif (PubMed:<a href="http://www.uniprot.org/citations/28991411" target="\_blank">28991411</a>, PubMed:<a href="http://www.uniprot.org/citations/32130408" target="\_blank">32130408</a>, PubMed:<a href="http://www.uniprot.org/citations/7604027" target="\_blank">7604027</a>, PubMed:<a href="http://www.uniprot.org/citations/7754385" target="\_blank">7754385</a>). Docking of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism (PubMed:<a href="http://www.uniprot.org/citations/7604027" target="\_blank">7604027</a>, PubMed:<a href="http://www.uniprot.org/citations/7754385" target="\_blank">7754385</a>). At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. Mediator of PR-DUB complex component BAP1 nuclear import; acts redundantly with KPNA1 and Transportin-1/TNPO1 (PubMed:<a href="http://www.uniprot.org/citations/35446349" target="\_blank">35446349</a>).

**Cellular Location**

Cytoplasm. Nucleus


**Tissue Location**


Expressed ubiquitously.

**Anti-KPNA2 Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-KPNA2 Monoclonal Antibody - Images**



Western blot analysis of KPNA2 expression in (1) HeLa cell lysate; (2) C6 cell lysate.