

Anti-alpha Internexin INA Monoclonal Antibody

Catalog # ABO14624

Specification

Anti-alpha Internexin INA Monoclonal Antibody - Product Information

Application WB, IHC, IF, ICC, IP, FC

Primary Accession

Host
Isotype

Q16352
Rabbit
Rabbit IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

Description

Anti-alpha Internexin INA Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-alpha Internexin INA Monoclonal Antibody - Additional Information

Gene ID 9118

Other Names

Alpha-internexin, Alpha-Inx, 66 kDa neurofilament protein, NF-66, Neurofilament-66, Neurofilament 5, INA, NEF5

Application Details

WB 1:500-1:2000
br>IHC 1:50-1:200
br>ICC/IF 1:50-1:200
br>IP 1:50
br>FC 1:20

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human alpha Internexin Class-IV neuronal intermediate filament that is able to self-assemble. It is involved in the morphogenesis of neurons. It may form an independent structural network without the involvement of other neurofilaments or it may cooperate with NF-L to form the filamentous backbone to which NF-M and NF-H attach to form the cross-bridges.

Purification

Affinity-chromatography

Storage Store at -20°C for one year. For short term


storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

Anti-alpha Internexin INA Monoclonal Antibody - Protein Information





Name INA

Synonyms NEF5

Function

Class-IV neuronal intermediate filament that is able to self- assemble. It is involved in the morphogenesis of neurons. It may form an independent structural network without the involvement of other neurofilaments or it may cooperate with NEFL to form the filamentous backbone to which NEFM and NEFH attach to form the cross-bridges. May also cooperate with the neuronal intermediate filament protein PRPH to form filamentous networks (By similarity).

Tissue Location


Found predominantly in adult CNS.

Anti-alpha Internexin INA Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-alpha Internexin INA Monoclonal Antibody - Images



Western blot analysis of alpha Internexin expression in SHSY5Y cell lysate.