

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody

Catalog # ABO14630

Specification

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody - Product Information

Application Primary Accession Host Isotype Reactivity Clonality Format **Description** Anti-RING2 / RING1B WB, IHC, IF, ICC, FC <u>099496</u> Rabbit Rabbit IgG Rat, Human, Mouse Monoclonal Liquid

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 6045

Other Names

E3 ubiquitin-protein ligase RING2, 2.3.2.27, Huntingtin-interacting protein 2-interacting protein 3, HIP2-interacting protein 3, Protein DinG, RING finger protein 1B, RING1b, RING finger protein 2, RING finger protein BAP-1, RING-type E3 ubiquitin transferase RING2, RNF2, BAP1, DING, HIPI3, RING1B

Application Details WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human RING2 / RING1B / RNF2 E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A, thereby playing a central role in histone code and gene regulation. H2A 'Lys-119' ubiquitination gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals.

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody - Protein Information



Name RNF2

Synonyms BAP1, DING, HIPI3, RING1B

Function

E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A (H2AK119Ub), thereby playing a central role in histone code and gene regulation (PubMed:15386022, PubMed:16359901, PubMed:21772249, PubMed:25355358, PubMed:25355358, PubMed:25519132, PubMed:25519132, PubMed:26151332, PubMed:26151332, PubMed:33864376, PubMed:33864376, PubMed:33864376, PubMed:33864376, PubMed:33864376, PubMed:16359901, PubMed:26151332). PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility (PubMed:26151332). E3 ubiquitin-protein ligase activity is enhanced by BMI1/PCGF4 (PubMed:21772249). Acts as the main E3 ubiquitin ligase on histone H2A of the PRC1 complex, while RING1 may rather act as a modulator of RNF2/RING2 activity (Probable). Association with the chromosomal DNA is cell-cycle dependent. In resting B- and T-lymphocytes, interaction with AURKB leads to block its activity, thereby maintaining transcription in resting lymphocytes (By similarity). Also acts as a negative regulator of autophagy by mediating ubiquitination of AMBRA1, leading to its subsequent degradation (By similarity).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q9CQJ4}. Chromosome {ECO:0000250|UniProtKB:Q9CQJ4}. Note=Enriched on inactive X chromosome (Xi) in female trophoblast stem (TS) cells as well as differentiating embryonic stem (ES) cells. The enrichment on Xi is transient during TS and ES cell differentiation. The association with Xi is mitotically stable in non-differentiated TS cells. Co-localizes with SAMD7 in nuclear polycomb bodies. {ECO:0000250|UniProtKB:Q9CQJ4}

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-RING2 / RING1B / RNF2 Rabbit Monoclonal Antibody - Images





Western blot analysis of RING2 / RING1B / RNF2 expression in HeLa cell lysate.