

# Anti-SEC23 Monoclonal Antibody

Catalog # ABO14640

## Specification

# Anti-SEC23 Monoclonal Antibody - Product Information

Application	WB, FC
Primary Accession	<u>Q15436</u>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	
Anti-SEC23 Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts	
with Human.	

# Anti-SEC23 Monoclonal Antibody - Additional Information

Gene ID 10484

**Other Names** Protein transport protein Sec23A, hSec23A, SEC23-related protein A, SEC23A (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=10701" target="\_blank">HGNC:10701</a>)

Application Details WB 1:1000-1:5000<br>FC 1:50

**Contents** Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human SEC23 Component of the COPII coat, that covers ER-derived vesicles involved in transport from the endoplasmic reticulum to the Golgi apparatus.

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

## Anti-SEC23 Monoclonal Antibody - Protein Information

Name SEC23A (HGNC:10701)



## Function

Component of the coat protein complex II (COPII) which promotes the formation of transport vesicles from the endoplasmic reticulum (ER). The coat has two main functions, the physical deformation of the endoplasmic reticulum membrane into vesicles and the selection of cargo molecules for their transport to the Golgi complex. Required for the translocation of insulin-induced glucose transporter SLC2A4/GLUT4 to the cell membrane (By similarity).

#### **Cellular Location**

Cytoplasmic vesicle, COPII-coated vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytosol. Note=Enriched at endoplasmic reticulum exit sites, also known as transitional endoplasmic reticulum (tER)

**Tissue Location** Ubiquitously expressed.

## **Anti-SEC23 Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### Anti-SEC23 Monoclonal Antibody - Images



Western blot analysis of SEC23 expression in HeLa cell lysate.