

Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody
Catalog # ABO14674

Specification

Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P10826
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody - Additional Information

Gene ID 5915

Other Names

Retinoic acid receptor beta, RAR-beta, HBV-activated protein, Nuclear receptor subfamily 1 group B member 2, RAR-epsilon, RARB, HAP, NR1B2

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Retinoic Acid Receptor beta Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes.

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody - Protein Information

Name RARB

Synonyms HAP, NR1B2**Function**

Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RXR/RAR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. In the absence or presence of hormone ligand, acts mainly as an activator of gene expression due to weak binding to corepressors (PubMed: [12554770](http://www.uniprot.org/citations/12554770)). The RXRA/RARB heterodimer can act as a repressor on the DR1 element and as an activator on the DR5 element (PubMed: [29021580](http://www.uniprot.org/citations/29021580)). In concert with RARG, required for skeletal growth, matrix homeostasis and growth plate function (By similarity).

Cellular Location

Nucleus. Cytoplasm [Isoform Beta-2]: Nucleus.


Tissue Location

Expressed in aortic endothelial cells (at protein level).


Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Retinoic Acid Receptor beta RARB Monoclonal Antibody - Images

Western blot analysis of Retinoic Acid Receptor beta expression in MCF7 cell lysate.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.