

Anti-EAAT3 Monoclonal Antibody
Catalog # ABO14713**Specification****Anti-EAAT3 Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	P43005
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-EAAT3 Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Anti-EAAT3 Monoclonal Antibody - Additional Information

Gene ID 6505

Other Names

Excitatory amino acid transporter 3, Excitatory amino-acid carrier 1, Neuronal and epithelial glutamate transporter, Sodium-dependent glutamate/aspartate transporter 3, Solute carrier family 1 member 1, SLC1A1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=10939)
HGNC:10939

Application Details

WB 1:1000-1:5000
IHC 1:100-1:500
ICC/IF 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human EAAT3 Transports L-glutamate and also L- and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium. Negatively regulated by ARL6IP5.

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-EAAT3 Monoclonal Antibody - Protein Information

Name SLC1A1 ([HGNC:10939](#))

Function

Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed:[21123949](http://www.uniprot.org/citations/21123949), PubMed:[26690923](http://www.uniprot.org/citations/26690923), PubMed:[33658209](http://www.uniprot.org/citations/33658209), PubMed:[7521911](http://www.uniprot.org/citations/7521911), PubMed:[7914198](http://www.uniprot.org/citations/7914198), PubMed:[8857541](http://www.uniprot.org/citations/8857541)). Can also transport L-cysteine (PubMed:[21123949](http://www.uniprot.org/citations/21123949)). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed:[26690923](http://www.uniprot.org/citations/26690923), PubMed:[33658209](http://www.uniprot.org/citations/33658209), PubMed:[7521911](http://www.uniprot.org/citations/7521911), PubMed:[8857541](http://www.uniprot.org/citations/8857541)). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed:[26690923](http://www.uniprot.org/citations/26690923), PubMed:[8857541](http://www.uniprot.org/citations/8857541)). Plays an important role in L- glutamate and L-aspartate reabsorption in renal tubuli (PubMed:[21123949](http://www.uniprot.org/citations/21123949)). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity). Contributes to glutathione biosynthesis and protection against oxidative stress via its role in L-glutamate and L-cysteine transport (By similarity). Negatively regulated by ARL6IP5 (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Apical cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Synapse, synaptosome {ECO:0000250|UniProtKB:P51906}. Early endosome membrane {ECO:0000250|UniProtKB:P51906}. Late endosome membrane {ECO:0000250|UniProtKB:P51906}. Recycling endosome membrane {ECO:0000250|UniProtKB:P51906}

Tissue Location


Expressed in all tissues tested including liver, muscle, testis, ovary, retinoblastoma cell line, neurons and brain (in which there was dense expression in substantia nigra, red nucleus, hippocampus and in cerebral cortical layers)

Anti-EAAT3 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EAAT3 Monoclonal Antibody - Images



Western blot analysis of EAAT3 expression in Human fetal brain lysate.