

Anti-PERP Monoclonal Antibody
Catalog # ABO14730**Specification**

Anti-PERP Monoclonal Antibody - Product Information

Application	WB, IF, ICC
Primary Accession	Q96FX8
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-PERP Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Anti-PERP Monoclonal Antibody - Additional Information

Gene ID 64065

Other Names

p53 apoptosis effector related to PMP-22 {ECO:0000312|HGNC:HGNC:17637},
Keratinocyte-associated protein 1, KCP-1, P53-induced protein PIGPC1 {ECO:0000303|Ref.3},
Transmembrane protein THW, PERP (<a
href="http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=17637"
target="_blank">HGNC:17637)

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human PERP

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-PERP Monoclonal Antibody - Protein Information

Name PERP ([HGNC:17637](#))

Function

Component of intercellular desmosome junctions (By similarity). Plays a role in stratified epithelial integrity and cell- cell adhesion by promoting desmosome assembly (By similarity). Thereby plays a role in barrier function of the skin against infection (By similarity). Plays a role in mammary epithelial tissue homeostasis and remodeling during and after pregnancy, potentially via its involvement in desmosome cell-cell junctions (By similarity). Required for tooth enamel development via facilitating desmosome-mediated ameloblast adhesion to the stratum intermedium during the transitional stage of amelogenesis (By similarity). May also play a role in downstream transcriptional regulation of other genes involved in amelogenesis such as AMBN, ENAM, MMP20 and KLK4 (By similarity). Plays a role as an effector in the TP53-dependent apoptotic pathway (By similarity). Positively regulates apoptosis in T-helper 17 (Th17) cell populations via caspase-dependent signaling (By similarity). Promotes neutrophil transepithelial migration in response to chemoattractants such as heparin A3 (HXA3), N-Formylmethionyl-leucyl-phenylalanine (fMLP) and CXCL8/IL-8 (PubMed:25486861). Required for neutrophil transepithelial migration in response to S.typhimurium infection (PubMed:25486861). May act as a positive regulator of endothelial cell apoptosis in response to blood flow-derived shear stress (By similarity).

Cellular Location

Cell junction, desmosome {ECO:0000250|UniProtKB:Q9JK95}. Cell membrane; Multi-pass membrane protein. Cytoplasm. Note=Associated with desmosomes (By similarity). Colocalizes with KRT14 in the cell membrane (PubMed:31898316). Clusters in a punctate pattern throughout the epithelial cytoplasm, in response to S.typhimurium infection (PubMed:25486861). {ECO:0000250|UniProtKB:Q9JK95, ECO:0000269|PubMed:25486861, ECO:0000269|PubMed:31898316}

Tissue Location


Expressed in skin, heart, placental, liver, pancreas, keratinocytes and dermal fibroblasts. May translocate to the intestinal apical epithelial cell surface via sipA and sctB1/sipC- promoted exocytic translocation following infection by S. Typhimurium (PubMed:25486861, PubMed:27078059).


Anti-PERP Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PERP Monoclonal Antibody - Images





Western blot analysis of PERP expression in A431 cell lysate.