

**Anti-ABHD5 Monoclonal Antibody**  
**Catalog # ABO14771****Specification**

---

**Anti-ABHD5 Monoclonal Antibody - Product Information**

Application	WB, IHC, FC
Primary Accession	<a href="#">Q8WTS1</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-ABHD5 Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human.

**Anti-ABHD5 Monoclonal Antibody - Additional Information**

**Gene ID** 51099

**Other Names**

1-acylglycerol-3-phosphate O-acyltransferase ABHD5, 2.3.1.51, Abhydrolase domain-containing protein 5, Lipid droplet-binding protein CGI-58, ABHD5 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=21396](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=21396)) target="\_blank">HGNC:21396</a>), NCIE2

**Application Details**

WB 1:500-1:2000<br>IHC 1:100-1:500<br>FC 1:100

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human ABHD5 Lysophosphatidic acid acyltransferase which functions in phosphatidic acid biosynthesis. May regulate the cellular storage of triacylglycerol through activation of the phospholipase PNPLA2. Involved in keratinocyte differentiation.

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-ABHD5 Monoclonal Antibody - Protein Information**

**Name** ABHD5 ([HGNC:21396](#))

**Synonyms** NCIE2

### Function

Coenzyme A-dependent lysophosphatidic acid acyltransferase that catalyzes the transfer of an acyl group on a lysophosphatidic acid (PubMed:<a href="http://www.uniprot.org/citations/18606822" target="\_blank">18606822</a>). Functions preferentially with 1-oleoyl- lysophosphatidic acid followed by 1-palmitoyl-lysophosphatidic acid, 1-stearoyl-lysophosphatidic acid and 1-arachidonoyl-lysophosphatidic acid as lipid acceptor. Functions preferentially with arachidonoyl-CoA followed by oleoyl-CoA as acyl group donors (By similarity). Functions in phosphatidic acid biosynthesis (PubMed:<a href="http://www.uniprot.org/citations/18606822" target="\_blank">18606822</a>). May regulate the cellular storage of triacylglycerol through activation of the phospholipase PNPLA2 (PubMed:<a href="http://www.uniprot.org/citations/16679289" target="\_blank">16679289</a>). Involved in keratinocyte differentiation (PubMed:<a href="http://www.uniprot.org/citations/18832586" target="\_blank">18832586</a>). Regulates lipid droplet fusion (By similarity).

### Cellular Location

Cytoplasm. Lipid droplet {ECO:0000250|UniProtKB:Q9DBL9}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9DBL9}. Note=Colocalized with PLIN and ADRP on the surface of lipid droplets. The localization is dependent upon the metabolic status of the adipocytes and the activity of PKA (By similarity).

### Tissue Location


Widely expressed in various tissues, including lymphocytes, liver, skeletal muscle and brain. Expressed by upper epidermal layers and dermal fibroblasts in skin, hepatocytes and neurons (at protein level).


## Anti-ABHD5 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-ABHD5 Monoclonal Antibody - Images





Western blot analysis of ABHD5 expression in HepG2 cell lysate.