

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9)
Catalog # ABO14833**Specification****Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) - Product Information**

Application	WB, IF, ICC, FC
Primary Accession	P20339
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) . Tested in Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500 µg/ml.

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) - Additional Information

Gene ID 5868

Other Names

Ras-related protein Rab-5A, 3.6.5.2, RAB5A, RAB5

Calculated MW

24 kDa KDa

Application Details

Western blot, 0.1-0.5 µg/ml
 Immunocytochemistry/Immunofluorescence, 2 µg/ml
 Flow Cytometry, 1-3 µg/1x10⁶ cells

Subcellular Localization

Early endosome membrane. Lipid-anchor. Endosome membrane. Cytosol. Cell membrane. Cytoplasmic side. Melanosome. Cytoplasmic vesicle. Ruffle. Membrane. Phagosome membrane.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Rab5, different from the related mouse and rat sequences by three amino acids.

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one

month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) - Protein Information

Name RAB5A ([HGNC:9783](#))

Synonyms RAB5

Function

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB5A is required for the fusion of plasma membranes and early endosomes (PubMed:10818110, PubMed:14617813, PubMed:15378032, PubMed:16410077). Contributes to the regulation of filopodia extension (PubMed:14978216). Required for the exosomal release of SDCBP, CD63, PDCD6IP and syndecan (PubMed:22660413). Regulates maturation of apoptotic cell-containing phagosomes, probably downstream of DYN2 and PIK3C3 (By similarity).

Cellular Location


Cell membrane; Lipid-anchor; Cytoplasmic side. Early endosome membrane; Lipid- anchor. Melanosome. Cytoplasmic vesicle. Cell projection, ruffle {ECO:0000250|UniProtKB:P18066}. Membrane Cytoplasm, cytosol. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q9CQD1}. Endosome membrane Note=Enriched in stage I melanosomes (PubMed:17081065). Alternates between membrane-bound and cytosolic forms (Probable) {ECO:0000269|PubMed:17081065, ECO:0000305}

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) - Images






Figure 1. IF analysis of RAB5A using anti-RAB5A antibody (M01891-1).

RAB5A was detected in immunocytochemical section of MCF7 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2 μ g/mL mouse anti-RAB5A Antibody (M01891-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 2. Western blot analysis of Rab5/RAB5A using anti-Rab5/RAB5A antibody (M01891-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

- Lane 1: human Hela whole cell lysates,
- Lane 2: human placenta tissue lysates,
- Lane 3: human A549 whole cell lysates.
- Lane 4: human A375 whole cell lysates,
- Lane 5: human SW620 whole cell lysates,
- Lane 6: human U20S whole cell lysates,
- Lane 7: human HEK293 whole cell lysates,
- Lane 8: human U87 whole cell lysates,
- Lane 9: rat brain tissue lysates,
- Lane 10: mouse brain tissue lysates,
- Lane 11: mouse Neuro-2a whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Rab5/RAB5A antigen affinity purified monoclonal antibody (Catalog #

M01891-1) at 0.5 $\mu\text{g/mL}$ overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Rab5/RAB5A at approximately 24KD. The expected band size for Rab5/RAB5A is at 24KD.




Figure 3. Flow Cytometry analysis of A431 cells using anti-Rab5/RAB5A antibody (M01891-1). Overlay histogram showing A431 cells stained with M01891-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Rab5/RAB5A Antibody (M01891-1, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Anti-Rab5/RAB5A Antibody Picoband™ (monoclonal, 3E9) - Background

RAB5A (Ras-associated protein RAB5A), also called RAB5, is a protein that in humans is encoded by the RAB5A gene. RAB5 is a rate-limiting component of the machinery regulating the kinetics of membrane traffic in the early endocytic pathway. The RAB5A gene is mapped on 3p24.3. RAB5 is indispensable for a form of receptor tyrosine kinase-induced actin remodeling called circular ruffling. It signals to the actin cytoskeleton through RNTRE, a RAB5-specific GTPase-activating protein (GAP). RAB5 activity on phagosome membranes began to increase on disassembly of the actin coat encapsulating phagosomes. In addition, RAB5 activation is either continuous or repetitive for up to 10 minutes, but it ends before the collapse of engulfed apoptotic cells. Expression of a dominant-negative mutant of RAB5 delayed this collapse of apoptotic thymocytes, showing a role for RAB5 in phagosome maturation.