

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5)
Catalog # ABO14948**Specification****Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q99757
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) . Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) - Additional Information

Gene ID 25828

Other Names

Thioredoxin, mitochondrial, MTRX, Mt-Trx, Thioredoxin-2, TXN2, TRX2

Calculated MW

14 kDa KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat
Immunocytochemistry/Immunofluorescence, 2 µg/ml, Human
Flow Cytometry, 1-3 µg/1x10⁶ cells, Human

Subcellular Localization

Mitochondrion.

Tissue Specificity

Widely expressed in adult (at protein level) and fetal tissues.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived human Thioredoxin 2/TXN2 recombinant protein (Position: T60-G166).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) - Protein Information

Name TXN2

Synonyms TRX2

Function

Important for the control of mitochondrial reactive oxygen species homeostasis, apoptosis regulation and cell viability (PubMed:12032145, PubMed:12080052, PubMed:26626369) Is involved in various redox reactions including the reduction of protein disulfide bonds, through the reversible oxidation of its active center dithiol to a disulfide (By similarity).

Cellular Location

Mitochondrion


Tissue Location

Widely expressed in adult (at protein level) and fetal tissues.

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) - Images




Figure 2. IF analysis of Thioredoxin 2/TXN2 using anti-Thioredoxin 2/TXN2 antibody (M04586-1). Thioredoxin 2/TXN2 was detected in immunocytochemical section of Hela cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2 $\mu\text{g}/\text{mL}$ mouse anti-Thioredoxin 2/TXN2 Antibody (M04586-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 3. Flow Cytometry analysis of HL-60 cells using anti-Thioredoxin 2/TXN2 antibody (M04586-1).

Overlay histogram showing HL-60 cells stained with M04586-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Thioredoxin 2/TXN2 Antibody (M04586-1, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 1. Western blot analysis of Thioredoxin 2/TXN2 using anti-Thioredoxin 2/TXN2 antibody (M04586-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates;

Lane 2: human Raji whole cell lysates;

Lane 3: human K562 whole cell lysates;

Lane 4: human A549 whole cell lysates;

Lane 5: human Caco-2 whole cell lysates;

Lane 6: human THP-1 whole cell lysates;

Lane 7: rat PC-12 whole cell lysates;

Lane 8: mouse RAW264.7 whole cell lysates

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Thioredoxin 2/TXN2 antigen affinity purified monoclonal antibody (Catalog # M04586-1) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for CTNNA1 at approximately 14KD. The expected band size for CTNNA1 is at 14KD.




Figure 4. IHC analysis of Thioredoxin 2/TXN2 using anti-Thioredoxin 2/TXN2 antibody (M04586-1).

Thioredoxin 2/TXN2 was detected in paraffin-embedded section of human gastric cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 µg/ml mouse anti-Thioredoxin 2/TXN2 Antibody (M04586-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 5. IHC analysis of Thioredoxin 2/TXN2 using anti-Thioredoxin 2/TXN2 antibody (M04586-1). Thioredoxin 2/TXN2 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 µg/ml mouse anti-Thioredoxin 2/TXN2 Antibody (M04586-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 6. IHC analysis of Thioredoxin 2/TXN2 using anti-Thioredoxin 2/TXN2 antibody (M04586-1). Thioredoxin 2/TXN2 was detected in paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 µg/ml mouse anti-Thioredoxin 2/TXN2 Antibody (M04586-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

Anti-Thioredoxin 2/TXN2 Antibody Picoband™ (monoclonal, 7B5) - Background

Thioredoxin, mitochondrial also known as thioredoxin-2 is a protein that in humans is encoded by the TXN2 gene on chromosome 22. It is mapped to 22q12.3. This nuclear gene encodes a mitochondrial member of the thioredoxin family, a group of small multifunctional redox-active proteins. The encoded protein may play important roles in the regulation of the mitochondrial membrane potential and in protection against oxidant-induced apoptosis.