

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5)
Catalog # ABO14995**Specification****Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) - Product Information**

Application	WB, IHC
Primary Accession	P00915
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) . Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) - Additional Information

Gene ID 759

Other Names

Carbonic anhydrase 1, 4.2.1.1, Carbonate dehydratase I, Carbonic anhydrase B, CAB, Carbonic anhydrase I, CA-I, Cyanamide hydratase CA1, 4.2.1.69, CA1

Calculated MW

29 kDa KDa

Application Details

Western blot, 0.25-0.5 µg/ml, Human, Mouse, Rat
 Immunohistochemistry (Paraffin-embedded Section), 2-5 µg/ml, Human

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na2HPO4.

Immunogen

E.coli-derived human CA1 recombinant protein (Position: D9-F261). Human CA1 shares 78.5% and 81% amino acid (aa) sequence identity with mouse and rat CA1, respectively.

Purification

Immunogen affinity purified.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored

frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) - Protein Information

Name CA1

Function

Catalyzes the reversible hydration of carbon dioxide (PubMed:10550681, PubMed:16506782, PubMed:16686544, PubMed:16807956, PubMed:17127057, PubMed:17314045, PubMed:17407288, PubMed:18618712, PubMed:19186056, PubMed:19206230). Can hydrate cyanamide to urea (PubMed:10550681).

Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:B0BNN3}.

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) - Images






Figure 1. Western blot analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human K562 whole cell lysates,

Lane 2: rat spleen tissue lysates,

Lane 3: mouse spleen tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Carbonic Anhydrase I/CA1 antigen affinity purified monoclonal antibody (Catalog # M00170-2) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Carbonic Anhydrase I/CA1 at approximately 29KD. The expected band size for Carbonic Anhydrase I/CA1 is at 29KD.




Figure 2. IHC analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Carbonic Anhydrase I/CA1 was detected in paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Carbonic Anhydrase I/CA1 Antibody (M00170-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 3. IHC analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Carbonic Anhydrase I/CA1 was detected in paraffin-embedded section of human gastric cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Carbonic Anhydrase I/CA1 Antibody (M00170-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IHC analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Carbonic Anhydrase I/CA1 was detected in paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Carbonic Anhydrase I/CA1 Antibody (M00170-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 5. IHC analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Carbonic Anhydrase I/CA1 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Carbonic Anhydrase I/CA1 Antibody (M00170-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30

minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 6. IHC analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Carbonic Anhydrase I/CA1 was detected in paraffin-embedded section of human pancreatic cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Carbonic Anhydrase I/CA1 Antibody (M00170-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 7. IHC analysis of Carbonic Anhydrase I/CA1 using anti-Carbonic Anhydrase I/CA1 antibody (M00170-2).

Carbonic Anhydrase I/CA1 was detected in paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Carbonic Anhydrase I/CA1 Antibody (M00170-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

Anti-Carbonic Anhydrase I/CA1 Picoband™ Antibody (monoclonal, 2B5) - Background

Carbonic anhydrase 1 is an enzyme that in humans is encoded by the CA1 gene. It is a member of the Carbonic anhydrase. The CA1 gene is mapped to 8q22. CAI has got about 260 amino acids. This protein is highly expressed in erythrocytes. As catalysts of the reversible hydration of carbon dioxide, CAI participates in a variety of biologic processes like respiration, calcification, acid-base balance etc.