

**Anti- IL-17F Monoclonal Antibody**  
**Catalog # ABO15021****Specification**

---

**Anti- IL-17F Monoclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q7TNI7</a>
Host	Mouse
Isotype	Mouse IgG1, κ
Reactivity	Mouse
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti- IL-17F Monoclonal Antibody . Tested in IHC-P applications. This antibody reacts with Mouse.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500 µ/ml.

**Anti- IL-17F Monoclonal Antibody - Additional Information**

**Gene ID** 257630

**Other Names**

Interleukin-17F, IL-17F, IL17f

**Application Details**

Immunohistochemistry (Paraffin-embedded Section), 2-5 µg/ml, Mouse<br>

**Protein Name**

Interleukin-17F

**Contents**

PBS, pH 7.0. Contains no stabilizers or preservatives

**Immunogen**

Mouse IL-17F

**Purification**

Immunogen affinity purified.

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti- IL-17F Monoclonal Antibody - Protein Information**

**Name** IL17f

## Function

Effector cytokine of innate and adaptive immune system involved in antimicrobial host defense and maintenance of tissue integrity (PubMed:<a href="http://www.uniprot.org/citations/18025225" target="\_blank">18025225</a>, PubMed:<a href="http://www.uniprot.org/citations/19144317" target="\_blank">19144317</a>, PubMed:<a href="http://www.uniprot.org/citations/23255360" target="\_blank">23255360</a>). IL17A- IL17F signals via IL17RA-IL17RC heterodimeric receptor complex, triggering homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter through SEFIR domains. This leads to downstream TRAF6- mediated activation of NF-kappa-B and MAPkinase pathways ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (PubMed:<a href="http://www.uniprot.org/citations/15477493" target="\_blank">15477493</a>, PubMed:<a href="http://www.uniprot.org/citations/17911633" target="\_blank">17911633</a>, PubMed:<a href="http://www.uniprot.org/citations/18025225" target="\_blank">18025225</a>). IL17A-IL17F is primarily involved in host defense against extracellular bacteria and fungi by inducing neutrophilic inflammation (PubMed:<a href="http://www.uniprot.org/citations/18025225" target="\_blank">18025225</a>, PubMed:<a href="http://www.uniprot.org/citations/23255360" target="\_blank">23255360</a>). As signature effector cytokine of T-helper 17 cells (Th17), primarily induces neutrophil activation and recruitment at infection and inflammatory sites (PubMed:<a href="http://www.uniprot.org/citations/18025225" target="\_blank">18025225</a>). Stimulates the production of antimicrobial beta- defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (PubMed:<a href="http://www.uniprot.org/citations/19144317" target="\_blank">19144317</a>). IL17F homodimer can signal via IL17RC homodimeric receptor complex, triggering downstream activation of TRAF6 and NF- kappa-B signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/28813677" target="\_blank">28813677</a>). Via IL17RC induces transcriptional activation of IL33, a potent cytokine that stimulates group 2 innate lymphoid cells and adaptive T-helper 2 cells involved in pulmonary allergic response to fungi (PubMed:<a href="http://www.uniprot.org/citations/28813677" target="\_blank">28813677</a>). Likely via IL17RC, promotes sympathetic innervation of peripheral organs by coordinating the communication between gamma-delta T cells and parenchymal cells. Stimulates sympathetic innervation of thermogenic adipose tissue by driving TGFB1 expression (PubMed:<a href="http://www.uniprot.org/citations/32076265" target="\_blank">32076265</a>). Regulates the composition of intestinal microbiota and immune tolerance by inducing antimicrobial proteins that specifically control the growth of commensal Firmicutes and Bacteroidetes (PubMed:<a href="http://www.uniprot.org/citations/29915298" target="\_blank">29915298</a>).

## Cellular Location

Secreted.

## Tissue Location

Expressed by T-helper 17 cells (Th17) (at protein level). The expression pattern reflects the differentiation state. In fully differentiated Th17 cells, IL17A-IL17F heterodimers are produced at higher levels than IL17A-IL17A and IL17F-IL17F dimers (PubMed:18025225). Dominantly secreted in intestine (PubMed:29915298) Expressed by resident cells of the lamina propria, both epithelial cells and immune cell subsets including natural killer cells, dendritic cells, macrophages and various T and B cell subsets (PubMed:16990136, PubMed:29915298). Expressed by epithelial cells and innate immune cells in the colon (PubMed:19144317). Expressed in group 3 innate lymphoid cells (PubMed:23255360, PubMed:29915298).

## Anti- IL-17F Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti- IL-17F Monoclonal Antibody - Images**




Figure 1. IHC analysis of IL-17F using anti-IL-17F antibody (M02062).

IL-17F was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 µg/ml mouse anti-IL-17F Antibody (M02062) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

#### **Anti- IL-17F Monoclonal Antibody - Background**

Interleukin 17F, also called IL17F is involved in the regulation of normal versus aberrant T-cell responses. This gene is mapped to 6p12.2. The protein encoded by this gene is a cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. It is suggested that targeting IL17 and IL17F or antagonizing IL17R might mitigate neutrophil-mediated inflammation in CF.