

Anti- IL 17A Monoclonal Antibody
Catalog # ABO15024**Specification**

Anti- IL 17A Monoclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q62386
Host	Mouse
Isotype	Mouse IgG1, κ
Reactivity	Mouse
Clonality	Monoclonal
Format	Lyophilized

Description

Anti- IL 17A Monoclonal Antibody . Tested in IHC-P applications. This antibody reacts with Mouse.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500 μ g/ml.

Anti- IL 17A Monoclonal Antibody - Additional Information

Gene ID 16171

Other Names

Interleukin-17A, IL-17, IL-17A, Cytotoxic T-lymphocyte-associated antigen 8, CTLA-8, Il17a, Ctl8, Il17

Application Details

Immunohistochemistry (Paraffin-embedded Section), 2-5 μ g/ml, Mouse

Protein Name

Interleukin-17A

Contents

PBS, pH 7.0. Contains no stabilizers or preservatives

Immunogen

Mouse IL-17A cross-linked to OVA

Purification

Immunogen affinity purified.

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti- IL 17A Monoclonal Antibody - Protein Information

Name Il17a**Synonyms** Ctlα8, Il17**Function**

Effector cytokine of innate and adaptive immune system involved in antimicrobial host defense and maintenance of tissue integrity (PubMed: [18025225](http://www.uniprot.org/citations/18025225) , PubMed: [19144317](http://www.uniprot.org/citations/19144317) , PubMed: [26431948](http://www.uniprot.org/citations/26431948)). Signals via IL17RA-IL17RC heterodimeric receptor complex, triggering homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter. This leads to downstream TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (PubMed: [16200068](http://www.uniprot.org/citations/16200068) , PubMed: [17911633](http://www.uniprot.org/citations/17911633) , PubMed: [19144317](http://www.uniprot.org/citations/19144317) , PubMed: [26431948](http://www.uniprot.org/citations/26431948)). Plays an important role in connecting T cell-mediated adaptive immunity and acute inflammatory response to destroy extracellular bacteria and fungi. As a signature effector cytokine of T-helper 17 cells (Th17), primarily induces neutrophil activation and recruitment at infection and inflammatory sites (PubMed: [18025225](http://www.uniprot.org/citations/18025225)). In airway epithelium, mediates neutrophil chemotaxis via induction of CXCL1 and CXCL5 chemokines (PubMed: [18025225](http://www.uniprot.org/citations/18025225) , PubMed: [27923703](http://www.uniprot.org/citations/27923703)). In secondary lymphoid organs, contributes to germinal center formation by regulating the chemotactic response of B cells to CXCL12 and CXCL13, enhancing retention of B cells within the germinal centers, B cell somatic hypermutation rate and selection toward plasma cells (PubMed: [18157131](http://www.uniprot.org/citations/18157131)). Effector cytokine of a subset of gamma-delta T cells that functions as part of an inflammatory circuit downstream IL1B, TLR2 and IL23A-IL12B to promote neutrophil recruitment for efficient bacterial clearance (PubMed: [17372004](http://www.uniprot.org/citations/17372004) , PubMed: [20364087](http://www.uniprot.org/citations/20364087) , PubMed: [28709803](http://www.uniprot.org/citations/28709803)). Effector cytokine of innate immune cells including invariant natural killer cell (iNKT) and group 3 innate lymphoid cells that mediate initial neutrophilic inflammation (PubMed: [17470641](http://www.uniprot.org/citations/17470641) , PubMed: [23255360](http://www.uniprot.org/citations/23255360)). Involved in the maintenance of the integrity of epithelial barriers during homeostasis and pathogen infection. Upon acute injury, has a direct role in epithelial barrier formation by regulating OCLN localization and tight junction biogenesis (PubMed: [26431948](http://www.uniprot.org/citations/26431948)). As part of the mucosal immune response induced by commensal bacteria, enhances host's ability to resist pathogenic bacterial and fungal infections by promoting neutrophil recruitment and antimicrobial peptides release (PubMed: [28709803](http://www.uniprot.org/citations/28709803)). In synergy with IL17F, mediates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (PubMed: [19144317](http://www.uniprot.org/citations/19144317)). Involved in antiviral host defense through various mechanisms (PubMed: [21946434](http://www.uniprot.org/citations/21946434) , PubMed: [26735852](http://www.uniprot.org/citations/26735852) , PubMed: [27795421](http://www.uniprot.org/citations/27795421)). Enhances immunity against West Nile virus by promoting T cell cytotoxicity (PubMed: [27795421](http://www.uniprot.org/citations/27795421)). May play a beneficial role in influenza A virus (H5N1) infection by enhancing B cell recruitment and immune response in the lung (PubMed: [21946434](http://www.uniprot.org/citations/21946434)).

target="_blank">21946434). Contributes to influenza A virus (H1N1) clearance by driving the differentiation of B-1a B cells, providing for production of virus-specific IgM antibodies at first line of host defense (PubMed:26735852).

Cellular Location

Secreted.

Tissue Location

Expressed by Th17 cell lineage (at protein level). The expression pattern reflects the differentiation state, with IL17A- IL17F heterodimers produced at higher levels than IL17A-IL17A and IL17F-IL17F dimers in fully differentiated Th17 cells (PubMed:16990136, PubMed:18025225). Expressed in innate lymphoid cells (at protein level) (PubMed:23255360, PubMed:28709803). Expressed in gamma-delta T cell subsets (at protein level) (PubMed:17372004, PubMed:20364087, PubMed:26431948, PubMed:28709803). Expressed in iNKT cells (at protein level) (PubMed:17470641).

Anti- IL 17A Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti- IL 17A Monoclonal Antibody - Images




Figure 1. IHC analysis of IL17A using anti-IL17A antibody (M00421-2).

IL17A was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 µg/ml mouse anti-IL17A Antibody (M00421-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as

secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

Anti- IL 17A Monoclonal Antibody - Background

Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.