

**Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4)**  
Catalog # ABO15047

**Specification**

**Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) - Product Information**

Application	WB
Primary Accession	<a href="#">Q9H9T3</a>
Host	Mouse
Isotype	Mouse IgG2a
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) . Tested in WB applications. This antibody reacts with Human, Mouse, Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) - Additional Information**

**Gene ID** 55140

**Other Names**

Elongator complex protein 3, hELP3, 2.3.1.311, ELP3 {ECO:0000303|PubMed:15902492, ECO:0000312|HGNC:HGNC:20696}

**Calculated MW**

62 kDa KDa

**Application Details**

Western blot, 0.1-0.25 µg/ml, Human, Mouse, Rat<br>

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na<sub>2</sub>HPO<sub>4</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human KAT9/ELP3, identical to the related mouse sequence.

**Purification**

Immunogen affinity purified.

**Storage**

**Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.**

## Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) - Protein Information

**Name** ELP3 {ECO:0000303|PubMed:15902492, ECO:0000312|HGNC:HGNC:20696}

### Function

Catalytic tRNA acetyltransferase subunit of the elongator complex which is required for multiple tRNA modifications, including mcm5U (5-methoxycarbonylmethyl uridine), mcm5s2U (5-methoxycarbonylmethyl-2-thiouridine), and ncm5U (5-carbamoylmethyl uridine) (PubMed:<a href="http://www.uniprot.org/citations/29415125" target="\_blank">29415125</a>). In the elongator complex, acts as a tRNA uridine(34) acetyltransferase by mediating formation of carboxymethyluridine in the wobble base at position 34 in tRNAs (By similarity). May also act as a protein lysine acetyltransferase by mediating acetylation of target proteins; such activity is however unclear in vivo and recent evidences suggest that ELP3 primarily acts as a tRNA acetyltransferase (PubMed:<a href="http://www.uniprot.org/citations/29415125" target="\_blank">29415125</a>). Involved in neurogenesis: regulates the migration and branching of projection neurons in the developing cerebral cortex, through a process depending on alpha-tubulin acetylation (PubMed:<a href="http://www.uniprot.org/citations/19185337" target="\_blank">19185337</a>). Required for acetylation of GJA1 in the developing cerebral cortex (By similarity).

### Cellular Location

Cytoplasm. Nucleus [Isoform 2]: Cytoplasm. Nucleus

### Tissue Location

Expressed in the cerebellum and spinal motor neurons.

## Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) - Images




Figure 1. Western blot analysis of KAT9/ELP3 using anti-KAT9/ELP3 antibody (M02833-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: human K562 whole cell lysates,  
Lane 2: human U20S whole cell lysates,  
Lane 3: human HepG2 whole cell lysates,  
Lane 4: human U87 whole cell lysates,  
Lane 5: rat lung tissue lysates,  
Lane 6: rat PC-12 whole cell lysates,  
Lane 7: mouse lung tissue lysates,  
Lane 8: mouse NIH/3T3 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-KAT9/ELP3 antigen affinity purified monoclonal antibody (Catalog # M02833-1) at 0.25 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for KAT9/ELP3 at approximately 62KD. The expected band size for KAT9/ELP3 is at 62KD.

#### **Anti-KAT9/ELP3 Antibody Picoband™ (monoclonal, 8B4) - Background**

Elongator complex protein 3, also named KAT9, is a protein that in humans is encoded by the ELP3 gene. It is mapped to 8p21.1. ELP3 is the catalytic histone acetyltransferase subunit of the RNA polymerase II elongator complex, which is a component of the RNA polymerase II (Pol II) holoenzyme and is involved in transcriptional elongation. ELP3 supports the migration and branching of projection neurons through acetylation of alpha-tubulin in the developing cerebral cortex. In mammals, ELP3 is important for paternal DNA demethylation after fertilization. ELP3 is potentially involved in cellular redox homeostasis by mediating the acetylation of glucose-6-phosphate dehydrogenase. Besides, ELP3 may play a role in chromatin remodeling and is involved in acetylation of histones H3 and probably H4.