

**Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3)**  
**Catalog # ABO15063****Specification****Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) - Product Information**

Application	WB, FC
Primary Accession	<a href="#">P07384</a>
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) . Tested in Flow Cytometry, WB applications. This antibody reacts with Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) - Additional Information**

**Gene ID** 823

**Other Names**

Calpain-1 catalytic subunit, 3.4.22.52, Calcium-activated neutral proteinase 1, CANP 1, Calpain mu-type, Calpain-1 large subunit, Cell proliferation-inducing gene 30 protein {ECO:0000303|Ref.3}, Micromolar-calpain, muCANP, CAPN1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=1476](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1476)), CANPL1

**Calculated MW**

82 kDa KDa

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human<br> Flow Cytometry, 1-3 µg/1x10<sup>6</sup> cells, Human<br>

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na<sub>2</sub>HPO<sub>4</sub>.

**Immunogen**

E.coli-derived human Calpain 1 recombinant protein (Position: Q396-A555). Human Calpain 1 shares 86% amino acid (aa) sequence identity with both mouse and rat Calpain 1.

**Purification**

Immunogen affinity purified.

**Storage**

**Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored**

**frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.**

## **Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) - Protein Information**

**Name** CAPN1 ([HGNC:1476](#))

**Synonyms** CANPL1

### **Function**

Calcium-regulated non-lysosomal thiol-protease which catalyzes limited proteolysis of substrates involved in cytoskeletal remodeling and signal transduction (PubMed:<a href="http://www.uniprot.org/citations/19617626" target="\_blank">19617626</a>, PubMed:<a href="http://www.uniprot.org/citations/21531719" target="\_blank">21531719</a>, PubMed:<a href="http://www.uniprot.org/citations/2400579" target="\_blank">2400579</a>). Proteolytically cleaves CTBP1 at 'Asn-375', 'Gly-387' and 'His-409' (PubMed:<a href="http://www.uniprot.org/citations/23707407" target="\_blank">23707407</a>). Cleaves and activates caspase-7 (CASP7) (PubMed:<a href="http://www.uniprot.org/citations/19617626" target="\_blank">19617626</a>).

### **Cellular Location**

Cytoplasm. Cell membrane. Note=Translocates to the plasma membrane upon Ca(2+) binding. In granular keratinocytes and in lower corneocytes, colocalizes with FLG and FLG2 (PubMed:21531719)

### **Tissue Location**


Ubiquitous.

## **Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) - Images**






Figure 1. Western blot analysis of Calpain 1 using anti-Calpain 1 antibody (M01943-4). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,  
Lane 2: human Hela whole cell lysates,  
Lane 3: human K562 whole cell lysates,  
Lane 4: human PC-3 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Calpain 1 antigen affinity purified monoclonal antibody (Catalog # M01943-4) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Calpain 1 at approximately 82KD. The expected band size for Calpain 1 is at 82KD.




Figure 2. Flow Cytometry analysis of A549 cells using anti-Calpain 1 antibody (M01943-4). Overlay histogram showing A549 cells stained with M01943-4 (Blue line).The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Calpain 1 Antibody (M01943-4, 1 µg/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 µg/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 µg/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

#### Anti-Calpain 1 Antibody Picoband™ (monoclonal, 2E3) - Background

CAPN1 is also known as CANP or muCL. The calpains, calcium-activated neutral proteases, are

nonlysosomal, intracellular cysteine proteases. The mammalian calpains include ubiquitous, stomach-specific, and muscle-specific proteins. The ubiquitous enzymes consist of heterodimers with distinct large, catalytic subunits associated with a common small, regulatory subunit. This gene encodes the large subunit of the ubiquitous enzyme, calpain 1. Several transcript variants encoding two different isoforms have been found for this gene.