

Anti-CHD3 Rabbit Monoclonal Antibody
Catalog # ABO15170**Specification**

Anti-CHD3 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q12873
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-CHD3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications.
This antibody reacts with Human, Mouse, Rat.

Anti-CHD3 Rabbit Monoclonal Antibody - Additional Information

Gene ID 1107

Other Names

Chromodomain-helicase-DNA-binding protein 3, CHD-3, 3.6.4.12, ATP-dependent helicase CHD3, Mi-2 autoantigen 240 kDa protein, Mi2-alpha, Zinc finger helicase, hZFH, CHD3

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CHD3

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-CHD3 Rabbit Monoclonal Antibody - Protein Information

Name CHD3

Function

ATP-dependent chromatin-remodeling factor that binds and distorts nucleosomal DNA (PubMed:<a

<http://www.uniprot.org/citations/28977666> target="_blank">28977666). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed:16428440, PubMed:28977666, PubMed:30397230, PubMed:9804427). Involved in transcriptional repression as part of the NuRD complex (PubMed:27068747). Required for anchoring centrosomal pericentrin in both interphase and mitosis, for spindle organization and centrosome integrity (PubMed:17626165).

Cellular Location

Nucleus, PML body. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome
Note=Associates with centrosomes in interphase and mitosis (PubMed:17626165). Localizes to sites of DNA damage (PubMed:28977666)

Tissue Location


Widely expressed.

Anti-CHD3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CHD3 Rabbit Monoclonal Antibody - Images



Western blot analysis of CHD3 expression in K562 cell lysate.