

Anti-p40 Rabbit Monoclonal Antibody
Catalog # ABO15311**Specification**

Anti-p40 Rabbit Monoclonal Antibody - Product Information

Application	WB, FC
Primary Accession	Q7Z6M1
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-p40 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human.

Anti-p40 Rabbit Monoclonal Antibody - Additional Information

Gene ID 10244

Other Names

Rab9 effector protein with kelch motifs, 40 kDa Rab9 effector protein, p40, RABEPK (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=16896), RAB9P40

Calculated MW

47 kDa KDa

Application Details

WB 1:500-1:2000
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human p40

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-p40 Rabbit Monoclonal Antibody - Protein Information

Name RABEPK ([HGNC:16896](#))

Synonyms RAB9P40

Function

Rab9 effector required for endosome to trans-Golgi network (TGN) transport.

Cellular Location


Cytoplasm. Endosome membrane. Note=Interaction with PIKFYVE and subsequent phosphorylation recruits it to the endosomal membrane

Anti-p40 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-p40 Rabbit Monoclonal Antibody - Images



Western blot analysis of p40 expression in HepG2 cell lysate.