

Anti-SAE1 Rabbit Monoclonal Antibody
Catalog # ABO15332**Specification**

Anti-SAE1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	Q9UBE0
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-SAE1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Rat.

Anti-SAE1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 10055

Other Names

SUMO-activating enzyme subunit 1, Ubiquitin-like 1-activating enzyme E1A, SUMO-activating enzyme subunit 1, N-terminally processed, SAE1, AOS1, SUA1, UBLE1A

Calculated MW

38 kDa KDa

Application Details

WB 1:1000-1:5000
ICC/IF 1:50-1:200
IP 1:50
FC 1:100

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human SAE1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-SAE1 Rabbit Monoclonal Antibody - Protein Information

Name SAE1

Synonyms AOS1, SUA1, UBLE1A

Function

The heterodimer acts as an E1 ligase for SUMO1, SUMO2, SUMO3, and probably SUMO4. It mediates ATP-dependent activation of SUMO proteins followed by formation of a thioester bond between a SUMO protein and a conserved active site cysteine residue on UBA2/SAE2.

Cellular Location

Nucleus.

Tissue Location

Expression level increases during S phase and drops in G2 phase (at protein level).

Anti-SAE1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SAE1 Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of SAE1 using anti-SAE1 antibody (M04753-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,

Lane 2: human Jurkat whole cell lysates,

Lane 3: human 293T whole cell lysates,

Lane 4: human MCF-7 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90

minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SAE1 antigen affinity purified monoclonal antibody (Catalog # M04753-2) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SAE1 at approximately 38 kDa. The expected band size for SAE1 is at 38 kDa.