

Anti-Jarid2 Rabbit Monoclonal Antibody
Catalog # ABO15566**Specification**

Anti-Jarid2 Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC
Primary Accession	Q92833
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Jarid2 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human.

Anti-Jarid2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 3720

Other Names

Protein Jumonji, Jumonji/ARID domain-containing protein 2, JARID2, JMJ

Calculated MW

139 kDa KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Jarid2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Jarid2 Rabbit Monoclonal Antibody - Protein Information

Name JARID2

Synonyms JMJ

Function

Regulator of histone methyltransferase complexes that plays an essential role in embryonic development, including heart and liver development, neural tube fusion process and hematopoiesis (PubMed:20075857). Acts as an accessory subunit for the core PRC2 (Polycomb repressive complex 2) complex, which mediates histone H3K27 (H3K27me3) trimethylation on chromatin (PubMed:20075857, PubMed:29499137, PubMed:31959557). Binds DNA and mediates the recruitment of the PRC2 complex to target genes in embryonic stem cells, thereby playing a key role in stem cell differentiation and normal embryonic development (PubMed:20075857). In cardiac cells, it is required to repress expression of cyclin-D1 (CCND1) by activating methylation of 'Lys-9' of histone H3 (H3K9me) by the GLP1/EHMT1 and G9a/EHMT2 histone methyltransferases (By similarity). Also acts as a transcriptional repressor of ANF via its interaction with GATA4 and NKX2-5 (By similarity). Participates in the negative regulation of cell proliferation signaling (By similarity). Does not have histone demethylase activity (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00355, ECO:0000255|PROSITE-ProRule:PRU00537, ECO:0000269|PubMed:20075857, ECO:0000269|PubMed:29499137}. Note=Colocalizes with the PRC2 complex on chromatin.

Tissue Location


During embryogenesis, predominantly expressed in neurons and particularly in dorsal root ganglion cells


Anti-Jarid2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Jarid2 Rabbit Monoclonal Antibody - Images





Western blot analysis of Jarid2 expression in NCCIT cell lysate.